

q7L
681
B16c
c.2
Birds

SMITHSONIAN
MUSEUM.

NORTH AMERICAN
BIRDS.

CATALOGUE

OF

NORTH AMERICAN BIRDS,

CHIEFLY IN THE MUSEUM OF THE

SMITHSONIAN INSTITUTION.

BY

SPENCER F. BAIRD,

ASSISTANT SECRETARY OF THE SMITHSONIAN INSTITUTION.



WASHINGTON:
SMITHSONIAN INSTITUTION.
OCTOBER, 1858.

VERSANDTUNG

1881

BRUNNEN

681
 B16c
 1859
 C. 2
 SCHHRB

ALPHABETICAL INDEX OF GENERA OF BIRDS.

(THE NUMBERS ARE THOSE OF THE FIRST SPECIES MENTIONED IN THE CATALOGUE.)

<p>A.</p> <p>Accipiter, 15 Actiturus, 545 Aetodromus, 531 Adamastor, 651 Aegialeus, 507 Aegialitis, 504 Aegiothus, 320 Aestrelata, 638 Agelaius, 401 Aix, 587 Alca, 710 Ammodromus, 341 Ampelis, 232 Anas, 576 Anorthura, 273 Anous, 696 Anser, 563 Anthus, 165 Antrostomus, 111 Aphriza, 511 Apobaptus, 732 Aquila, 39 Aramus, 480 Ardea, 487 Ardetta, 491 Archibuteo, 30 Ardenna, 647 Arquatella, 528 Astragalinus, 313 Astur, 14 Asturina, 33 Athene, 58 Atthis, 105 Audubonia, 489 Aythya, 591</p>	<p>Bucephala, 593 Buteo, 18 Butorides, 493</p> <p style="text-align: center;">C.</p> <p>Calamospiza, 377 Calidris, 534 Callipepla, 476 Campephilus, 72 Campylorhynchus, 600 Cardellina, 262 Cardinalis, 216 Carpodacus, 390 Cathartes, 305 Cathartes, 1 Catherpes, 263 Centrocercus, 462 Centronyx, 331 Centropneustes, 326 Centurus, 91 Cerorhinca, 717 Certhia, 275 Certhiola, 301 Coryle, 117 Chetura, 109 Chamaea, 274 Champepelia, 453 Charadrius, 503 Chauleclasmus, 584 Chen, 563 Chenalopez, 710 Chloephaga, 573 Chloroceryle, 118 Chondestes, 344 Chordeiles, 114 Chroicocephalus, 667 Chrysomitris, 310 Ciceronia, 722 Circus, 38 Cistothorus, 268 Coccyzus, 69 Colaptes, 99 Collyrio, 236 Columba, 445 Colymbus, 698 Contopus, 137 Conurus, 63</p>	<p>Corvus, 423 Coturnicops, 557 Coturniculus, 338 Cotyle, 229 Craxirex, 46 Creagrus, 679 Creiscus, 556 Crex, 558 Crotophaga, 66 Cupidonia, 464 Curvirostra, 318 Cyanocitta, 457 Cyanospiza, 383 Cyanura, 434 Cygnus, 561 Cyrtonyx, 477 Cyrtopelicanus, 615</p> <p style="text-align: center;">D.</p> <p>Dafila, 578 Daption, 639 Demigreggata, 482 Dendrocoryna, 574 Dendroica, 189 Diomedea, 630 Dolichonyx, 399 Dysporus, 399</p> <p style="text-align: center;">E.</p> <p>Ectopistes, 448 Elanus, 35 Embernagra, 373 Empidonax, 140 Eremophila, 302 Eremetes, 535 Eristatura, 609 Erolia, 529 Euphonia, 224 Euspiza, 378</p> <p style="text-align: center;">F.</p> <p>Falcinellus, 500 Falco, 5 Florida, 490 Fratercula, 713 Fregetta, 646</p>	<p>Fulica, 559 Fulix, 588 Fulmarus, 635</p> <p style="text-align: center;">G.</p> <p>Gallinago, 523 Gallinula, 560 Gambetta, 539 Garzetta, 485 Gennaia, 10 Geococcyx, 68 Geothlypis, 170 Glaucidium, 60 Glottis, 538 Goniaphea, 380 Graculus, 620 Grus, 478 Guiraca, 380 Gymnokitta, 431</p> <p style="text-align: center;">H.</p> <p>Haematopus, 512 Haliaetus, 40 Harelda, 597 Harporhynchus, 256 Helminthophaga, 180 Helmintherus, 178 Helospiza, 368 Herodias, 486 Hesperiphona, 303 Heteroscelus, 543 Hierofalco, 11 Himantopus, 518 Hirundo, 225 Histrionicus, 596 Hydrobata, 164 Hydrochelidon, 695 Hylotomus, 90 Hypotriorchis, 7</p> <p style="text-align: center;">I.</p> <p>Ibis, 498 Icteria, 176 Icterus, 408 Ictinia, 36 Ixoreus, 156</p>
---	---	--	--

Order II. SCANSORES. (Page 65.)

Family.	Page.	Sub-family.	Page.	Section.	Page.	Genus.	Page.	Sub-genus.	Page.	Species examined and identified.	Species not examined nor identified.	Extra limited species.	Current number of first species mentioned.
4. Psittacidae	66					29. Conurus, Kuhl	66			1			63
						Rhynchopsitta, Bon.	66					1	64
5. Trogonidae	66					30. Trogon, Moehr.				1			65
6. Cuculidae	71					31. Crotophaga, Linn.	71			2			66
						32. Geococcyx, Wag.	73			1			68
						33. Coccyzus, Vieill.	75			3			69
7. Picidae	79	Picinae	79	Piceae	80	34. Campephilus, Gray.	81			1		1	72
						35. Picus, Linn.	83	Trichopicus, Bp	83	4			74
								Dyctopicus, Bp.	83	2			78
								Phrenopicus, Bp.	83	1			80
								Xenopicus, Bd.	83	1			81
						36. Picoides, Lac.	97			3			82
						37. Sphyrapicus, Bd.	101			5			85
						38. Hylatomus, Baird ..	107			1			90
				Centureae	108	39. Centurus, Sw.	108			3			91
						40. Melanerpes, Sw.	112			3			94
				Colaptea	117	41. Colaptes, Sw.	117			3			99

Order III. INSESSORES. (Page 126.)

Sub-order A. STRISORES. (Page 128.)

10. Trochilidae	129					42. Lampornis, Sw.	130			1			100
						43. Trochilus, L.	131			2			101
						44. Sialaporphus, Sw.	133			2			103
						45. Atthis, Reich.	136			2			105
11. Cyscelidae	140					46. Panyptila, Cab.	140			1			107
						47. Nephocetes, Bd.	142			1			105
						48. Chaetura, Steph.	144			2			109
12. Caprimulgidae ..	146	Caprimulginae ..	146			49. Antrostomus, Gould.	146			3			111
						50. Chordeiles, Sw.	150			3			114

Sub-order B. CLAMATORES. (Page 156.)

13. Alcedinidae	157					51. Ceryle, Boie	157	Megaceryle, Rich.	157	1			117
								Chloroceryle, Kaup ..	157	1			118
14. Prionitidae	161					52. Momotus, Lath.	161			1			119
15. Colopterae	163	Psarinae	164			53. Pachyramphus, Gr.	164			1			120
						54. Bathmidurus, Cab.	165			1			121
		Tyranninae	166	Tyranni	167	55. Milvulus, Sw.	167			2			122
						56. Tyrannus, Cuv.	170			5		1	124
				Tyrannuli	167	57. Myiarchus, Cab.	177			4			130
						58. Sayornis, Bp.	182			3			134
						59. Contopus, Cab.	186			3			137
						60. Empidonax, Cab.	191			7			140
						61. Pyrocephalus, Gould	201			1			147

TABLE OF THE HIGHER GROUPS.

Sub-order C. OSCINES—Continued.

Family.	Page.	Sub-family.	Page.	Section.	Page.	Genus.	Page.	Sub-genus.	Page.	Species examined and identified.	Species not examined nor identified.	Extra limited species.	Current number of first species mentioned.
24. Dacnidae	App	Dacninae	App			109. Certhiola, Sand.	App			1			301
25. Alaudidae	402					110. Eremophila, Boie.	402			1			302
26. Fringillidae	406	Coccothraustinae	406			111. Hesperiphona, Bon.	408			1			303
						112. Pinalcola, Vieill.	409			1			304
						113. Carpodacus, Kaup.	411			4		1	305
						114. Chrysomitris, Boie.	418	Chrysomitris, Boie	418	1		2	310
								Astragalinus, Cab.	419	5			313
						115. Curvirostra, Scop.	436			2			318
						116. Aegiothus, Cab.	438			1	1		330
						117. Leucosticte, Sw.	430			1	2		322
						118. Plectrophanes, Mey.	431	Plectrophanes, Meyer.		1			325
								Centrophanes, Kaup.		4			336
		Spizellinae	436					Rhynchophanes, Bd.		1			330
						119. Centronyx, Baird.	440			1			331
						120. Passerculus, Bon.	441			5			332
						121. Poocetes, Baird.	447			1			337
						122. Coturniculus, Bon.	449			2	1		338
						123. Ammodromus, Sw.	452			3			341
						124. Chondestes, Sw.	455			1			344
						125. Zonotrichia, Sw.	457			5			345
						126. Junco, Wagler.	464			4		1	350
						127. Poospiza, Cab.	469			2			355
						128. Spizella, Bon.	471			6			357
						129. Melospiza, Bd.	476	Melospiza, Bd.	476	5			363
								Helospiza, Bd.	477	2			368
						130. Peucaea, Aud.	484			3			370
						131. Eubemagra, Lesq.	487			1			373
		Passercellinae	487			132. Passercella, Sw.	488			3			374
		Spizinae	490			133. Calanospiza, Bon.	492			1			377
						134. Euspiza, Bon.	493			2			378
						135. Guiraca, Sw.	496	Goniapheca, Bowd.	497	2			380
								Guiraca, Sw.	497	1			382
						136. Cyanospiza, Bd.	500			5			383
						137. Spermophila, Sw.	506			1			388
						138. Pyrrhuloxia, Bon.	507			1			389
						139. Cardinalis, Bon.	508			1			390
						140. Pipilo, Vieill.	510			8			391
27. Icteridae	521	Ageiainae	521			141. Dolichonyx, Sw.	522			1			399
						142. Molothrus, Sw.	523			1			400
						143. Agelaius, Vieill.	525			3			401
						144. Xanthocephalus, B.	531			1			404
						(145.) Trupialis, Bon.	533			7		(?) 1	405
						146. Sturnella, Vieill.	535			2			406
		Icterinae	540			147. Icterus, Daud.	540			8		1	408
		Quiscalinae	550			148. Scolopogon, Sw.	551			2			417
						149. Quiscalus, Vieill.	553			4			419
28. Corvidae	558	Corvinae	558			150. Corvus, Linn.	559			7			423
						151. Picocorvus, Bon.	572			1			430
						152. Gymnokitta, Max.	574			1			431
		Garrulinae	575			153. Pica, Briss.	576			2			432
						154. Cyanura, Sw.	579			3			434
						155. Cyanocitta, Str.	584			5			437
						156. Xanthoura, Bp.	589			1			442
						157. Perisoreus, Bon.	590			1			443
						158. Psithyrus, Rupp.	591			1			444

Order IV. RASORES. (Page 593.)

Sub-Order A. COLUMBAE. (Page 595.)

Family.	Page.	Sub-family.	Page.	Section.	Page.	Genus.	Page.	Sub-genus.	Page.	Species examined and identified.	Species not examined nor identified.	Extra limited species.	Current number of list species mentioned.
29. Columbidae.....	595	Columbinae.....	596	159. Columba, L.....	596	Columba, L.....	596	2	445	
						160. Ectopistes, Sw....	599	Patagioenas, Reich..	596	1	447	
		Zenaidinae.....	600	Zenaidae.....	600	161. Zenaida, Bon....	601	1	448	
						162. Melopelia, Bon....	602	1	449	
						163. Zenaidura, Bon....	603	1	450	
				Chamaepeliæ	164. Scardafella, Bon....	605	1	452	
						165. Chamaepelia, Sw....	606	1	453	
				Starnoenædæ.....	166. Oreopelia, Reich.	607	1	454	
						167. Starnoenas, Bon....	608	1	455	

Sub-Order B. GALLINAE. (Page 609.)

30. Penelopidae.....	610	168. Ortalida, Merrem..	610	1	456	
31. Phasianidae.....	613	Meleagrinae.....	613	169. Meleagris, Linn....	613	2	457	
32. Tetraonidae.....	619	170. Tetrao, Linn.....	630	3	459	
						171. Centrocercus, Sw....	634	1	462	
						172. Pedioecetes, Bd....	635	1	463	
						173. Cupidonia, Reich..	637	1	464	
						174. Bonasa, Steph.....	639	2	465	
						175. Lagopus, Vieill....	632	3	1	467	
33. Perdiciidae.....	638	Ortyginae.....	638	176. Ortyx, Steph.....	639	2	471	
						177. Oreortyx, Baird....	642	1	473	
						178. Lophortyx, Bon....	643	2	474	
						179. Callipepla, Wagler..	646	1	476	
						180. Cyrtonyx, Gould..	647	1	477	

Order V. GRALLATORES. (Page 650.)

Sub-Order A. HERODIONES. (Page 651.)

34. Gruidæ.....	658	181. Grus, Linn.....	653	3	478	
35. Aramidæ.....	657	182. Aramus, Vieill....	657	1	480	
36. Ardeidæ.....	659	Ardeinae.....	659	Ardeæ.....	659	183. Demigretta, Blyth.	660	3	482	
						184. Garzetta, Bon....	664	1	485	
						185. Herodias, Boie....	666	1	486	
						186. Ardea, Linn.....	667	2	487	
						187. Audubonia, Bp....	670	7	459	
						188. Florida, Bd.....	671	1	490	
						189. Ardetta, Gray....	672	1	491	
						190. Botaurus, Steph..	674	1	492	
						191. Eutorides, Blyth...	676	1	493	
						192. Nyctiardea, Sw....	678	1	495	
						193. Nyctherodias, Reic.	679	1	496	
37. Tantalidæ.....	681	Tantalinae.....	681	194. Tantalus, Linn....	682	1	497	
		Ibinae.....	681	195. Ibis, Mochr.....	682	Ibis, Mochr.....	2	498	
						Falcinellus, Bon....	1	500	
38. Plataleidæ.....	686	196. Platalea, L.....	686	1	501	
39. Phœnicopteridæ	687	197. Phœnicopterus, L.	687	1	502	

Geo. N. Laurence

Mar 1863

✓ 1	✓ 51	✓ 101	✓ 151	✓ 201	✓ 251	✓ 301	✓ 351
✓ 2	✓ 52	✓ 102	✓ 152	✓ 202	✓ 252	✓ 302	✓ 352
✓ 3	✓ 53	✓ 103	✓ 153	✓ 203	✓ 253	✓ 303	✓ 353
✓ 4	✓ 54	✓ 104	✓ 154 ^{hour}	✓ 204	✓ 254 ²⁵³	✓ 304	✓ 354
✓ 5	✓ 55	✓ 105	✓ 155	✓ 205	✓ 255	✓ 305	✓ 355
✓ 6	✓ 56	✓ 106	✓ 156	✓ 206	✓ 256	✓ 306	✓ 356
✓ 7	✓ 57	✓ 107	✓ 157	✓ 207	✓ 257	✓ 307	✓ 357
✓ 8	✓ 58	✓ 108	✓ 158	✓ 208	✓ 258	✓ 308	✓ 358
✓ 9	✓ 59	✓ 109	✓ 159	✓ 209	✓ 259	✓ 309	✓ 359
✓ 10	✓ 60 ^{hour}	✓ 110	✓ 160	✓ 210	✓ 260 ²⁵⁹	✓ 310	✓ 360
✓ 11	✓ 61	✓ 111	✓ 161	✓ 211	✓ 261	✓ 311	✓ 361
✓ 12	✓ 62	✓ 112	✓ 162	✓ 212	✓ 262 ²⁶¹	✓ 312	✓ 362
✓ 13	✓ 63	✓ 113	✓ 163	✓ 213	✓ 263	✓ 313	✓ 363
✓ 14	✓ 64	✓ 114	✓ 164	✓ 214	✓ 264	✓ 314	✓ 364
✓ 15	✓ 65	✓ 115	✓ 165	✓ 215	✓ 265	✓ 315	✓ 365
✓ 16	✓ 66	✓ 116	✓ 166	✓ 216	✓ 266	✓ 316	✓ 366
✓ 17	✓ 67	✓ 117	✓ 167	✓ 217	✓ 267	✓ 317	✓ 367
✓ 18	✓ 68	✓ 118	✓ 168	✓ 218	✓ 268	✓ 318	✓ 368
✓ 19	✓ 69	✓ 119	✓ 169	✓ 219	✓ 269	✓ 319	✓ 369
✓ 20	✓ 70	✓ 120	✓ 170	✓ 220	✓ 270	✓ 320	✓ 370
✓ 21	✓ 71	✓ 121	✓ 171	✓ 221	✓ 271	✓ 321	✓ 371
✓ 22	✓ 72	✓ 122	✓ 172	✓ 222	✓ 272	✓ 322	✓ 372
✓ 23	✓ 73	✓ 123	✓ 173	✓ 223	✓ 273	✓ 323	✓ 373
✓ 24	✓ 74	✓ 124	✓ 174	✓ 224	✓ 274	✓ 324	✓ 374
✓ 25	✓ 75	✓ 125	✓ 175	✓ 225	✓ 275	✓ 325	✓ 375
✓ 26	✓ 76	✓ 126	✓ 176	✓ 226	✓ 276	✓ 326	✓ 376
✓ 27	✓ 77	✓ 127	✓ 177	✓ 227	✓ 277	✓ 327	✓ 377
✓ 28	✓ 78	✓ 128	✓ 178	✓ 228	✓ 278	✓ 328	✓ 378
✓ 29	✓ 79	✓ 129	✓ 179	✓ 229	✓ 279	✓ 329	✓ 379
✓ 30	✓ 80	✓ 130	✓ 180	✓ 230	✓ 280	✓ 330	✓ 380
✓ 31	✓ 81	✓ 131	✓ 181	✓ 231	✓ 281	✓ 331	✓ 381
✓ 32	✓ 82	✓ 132	✓ 182	✓ 232 ²³¹	✓ 282	✓ 332	✓ 382
✓ 33	✓ 83	✓ 133	✓ 183	✓ 233	✓ 283	✓ 333	✓ 383
✓ 34	✓ 84	✓ 134	✓ 184	✓ 234	✓ 284	✓ 334	✓ 384
✓ 35	✓ 85	✓ 135	✓ 185	✓ 235	✓ 285	✓ 335	✓ 385
✓ 36	✓ 86	✓ 136	✓ 186	✓ 236	✓ 286	✓ 336	✓ 386
✓ 37	✓ 87	✓ 137	✓ 187	✓ 237	✓ 287	✓ 337	✓ 387
✓ 38	✓ 88	✓ 138	✓ 188	✓ 238	✓ 288	✓ 338	✓ 388
✓ 39	✓ 89	✓ 139	✓ 189	✓ 239	✓ 289	✓ 339	✓ 389
✓ 40	✓ 90	✓ 140	✓ 190	✓ 240	✓ 290 ²⁸⁹	✓ 340	✓ 390
✓ 41	✓ 91	✓ 141	✓ 191	✓ 241	✓ 291	✓ 341	✓ 391
✓ 42	✓ 92	✓ 142	✓ 192	✓ 242	✓ 292	✓ 342	✓ 392
✓ 43	✓ 93	✓ 143	✓ 193	✓ 243	✓ 293	✓ 343	✓ 393
✓ 44	✓ 94	✓ 144	✓ 194	✓ 244	✓ 294	✓ 344	✓ 394
✓ 45	✓ 95	✓ 145 ^{hour}	✓ 195	✓ 245	✓ 295	✓ 345	✓ 395
✓ 46	✓ 96	✓ 146	✓ 196	✓ 246	✓ 296	✓ 346	✓ 396
✓ 47	✓ 97	✓ 147	✓ 197	✓ 247	✓ 297	✓ 347	✓ 397
✓ 48	✓ 98	✓ 148	✓ 198	✓ 248	✓ 298	✓ 348	✓ 398
✓ 49	✓ 99	✓ 149	✓ 199	✓ 249	✓ 299	✓ 349	✓ 399
✓ 50	✓ 100	✓ 150 ^{hour}	✓ 200	✓ 250	✓ 300	✓ 350	✓ 400

✓ 401	✓ 451	✓ 501	✓ 551	✓ 601	✓ 651	✓ 701	751
✓ 402	✓ 452	✓ 502	✓ 552	✓ 602	✓ 652	✓ 702	752
✓ 403	✓ 453	✓ 503	✓ 553	✓ 603	✓ 653	✓ 703	753
✓ 404	✓ 454	✓ 504	✓ 554	✓ 604	✓ 654	✓ 704	754
✓ 405	✓ 455	✓ 505	✓ 555	✓ 605	✓ 655	✓ 705	755
✓ 406	✓ 456	✓ 506	✓ 556	✓ 606	✓ 656	✓ 706	756
✓ 407	✓ 457	✓ 507	✓ 557	✓ 607	✓ 657	✓ 707	757
✓ 408	✓ 458	✓ 508	✓ 558	✓ 608	✓ 658	✓ 708	758
✓ 409	✓ 459	✓ 509	✓ 559	✓ 609	✓ 659	✓ 709	759
✓ 410	✓ 460	✓ 510	✓ 560	✓ 610	✓ 660	✓ 710	760
✓ 411	✓ 461	✓ 511	✓ 561	✓ 611	✓ 661	✓ 711	761
✓ 412	✓ 462	✓ 512	✓ 562	✓ 612	✓ 662	✓ 712	762
✓ 413	✓ 463	✓ 513	✓ 563	✓ 613	✓ 663	✓ 713	763
✓ 414	✓ 464	✓ 514	✓ 564	✓ 614	✓ 664	✓ 714	764
✓ 415	✓ 465	✓ 515	✓ 565	✓ 615	✓ 665	✓ 715	765
✓ 416	✓ 466	✓ 516	✓ 566	✓ 616	✓ 666	✓ 716	766
✓ 417	✓ 467	✓ 517	✓ 567	✓ 617	✓ 667	✓ 717	767
✓ 418	✓ 468	✓ 518	✓ 568	✓ 618	✓ 668	✓ 718	768
✓ 419	✓ 469	✓ 519	✓ 569	✓ 619	✓ 669	✓ 719	769
✓ 420	✓ 470	✓ 520	✓ 570	✓ 620	✓ 670	✓ 720	770
✓ 421	✓ 471	✓ 521	✓ 571	✓ 621	✓ 671	✓ 721	771
✓ 422	✓ 472	✓ 522	✓ 572	✓ 622	✓ 672	✓ 722	772
✓ 423	✓ 473	✓ 523	✓ 573	✓ 623	✓ 673	✓ 723	773
✓ 424	✓ 474	✓ 524	✓ 574	✓ 624	✓ 674	✓ 724	774
✓ 425	✓ 475	✓ 525	✓ 575	✓ 625	✓ 675	✓ 725	775
✓ 426	✓ 476	✓ 526	✓ 576	✓ 626	✓ 676	✓ 726	776
✓ 427	✓ 477	✓ 527	✓ 577	✓ 627	✓ 677	✓ 727	777
✓ 428	✓ 478	✓ 528	✓ 578	✓ 628	✓ 678	✓ 728	778
✓ 429	✓ 479	✓ 529	✓ 579	✓ 629	✓ 679	✓ 729	779
✓ 430	✓ 480	✓ 530	✓ 580	✓ 630	✓ 680	✓ 730	780
✓ 431	✓ 481	✓ 531	✓ 581	✓ 631	✓ 681	✓ 731	781
✓ 432	✓ 482	✓ 532	✓ 582	✓ 632	✓ 682	✓ 732	782
✓ 433	✓ 483	✓ 533	✓ 583	✓ 633	✓ 683	✓ 733	783
✓ 434	✓ 484	✓ 534	✓ 584	✓ 634	✓ 684	✓ 734	784
✓ 435	✓ 485	✓ 535	✓ 585	✓ 635	✓ 685	✓ 735	785
✓ 436	✓ 486	✓ 536	✓ 586	✓ 636	✓ 686	✓ 736	786
✓ 437	✓ 487	✓ 537	✓ 587	✓ 637	✓ 687	✓ 737	787
✓ 438	✓ 488	✓ 538	✓ 588	✓ 638	✓ 688	✓ 738	788
✓ 439	✓ 489	✓ 539	✓ 589	✓ 639	✓ 689	✓ 739	789
✓ 440	✓ 490	✓ 540	✓ 590	✓ 640	✓ 690	✓ 740	790
✓ 441	✓ 491	✓ 541	✓ 591	✓ 641	✓ 691	✓ 741	791
✓ 442	✓ 492	✓ 542	✓ 592	✓ 642	✓ 692	✓ 742	792
✓ 443	✓ 493	✓ 543	✓ 593	✓ 643	✓ 693	✓ 743	793
✓ 444	✓ 494	✓ 544	✓ 594	✓ 644	✓ 694	✓ 744	794
✓ 445	✓ 495	✓ 545	✓ 595	✓ 645	✓ 695	✓ 745	795
✓ 446	✓ 496	✓ 546	✓ 596	✓ 646	✓ 696	✓ 746	796
✓ 447	✓ 497	✓ 547	✓ 597	✓ 647	✓ 697	✓ 747	797
✓ 448	✓ 498	✓ 548	✓ 598	✓ 648	✓ 698	✓ 748	798
✓ 449	✓ 499	✓ 549	✓ 599	✓ 649	✓ 699	✓ 749	799
✓ 450	✓ 500	✓ 550	✓ 600	✓ 650	✓ 700	✓ 750	800

TABLE OF THE HIGHER GROUPS.

XXIII

Order VI. NATATORES. (Page 752.)

Sub-order A. ANSERES. (Page 753.)

Family.	Sub-family.	Section.	Genus.	Sub-genus.	Page.	Species examined and identified.	Species not examined nor identified.	Extra limited species.	Current number of first species mentioned.		
46. Anatidae.....	Cygnae.....	757	230. Cygnus, Linn.....	757	Olor, Wagler.....	757	2	561		
		759	231. Anser, Linn.....	759	Chen, Boie.....	757	2	563		
	Anserinae.....	Anserae.....	759	232. Bernicla, Steph....	763	Anser, Linn.....	757	2	565	
			759	233. Chloephaga, Eyton.	769	Leucobepharon, Bd.	763	3	567	
			759	234. Dendrocygna, Sw..	769	Bernicla, Steph.....	763	2	570	
			759	235. Anas, Linn.....	773	Leucopareia, Rch....	763	1	572	
			759	236. Dafila, Leach....	774	573
			759	237. Nettion, Kaup....	777	574
			759	238. Querquedula, Steph.	779	576
			759	239. Spatula, Boie.....	781	578
			759	240. Chaulelasmus, Gray	782	579
			759	241. Mareca, Steph....	783	581
	Fuligulinae.....	786	242. Aix, Boie.....	785	583	
			786	243. Fulix, Sund.....	790	584	
			786	244. Aythya, Boie.....	793	585	
			786	245. Bucephala, Bd....	795	587	
			786	246. Histrionicus, Less.	798	588	
			786	247. Harelda, Leach....	800	591	
			786	248. Polysticta, Eyton..	801	593	
			786	249. Lamprolitta, Brit.	802	596	
			786	250. Campylaeus, Gr....	803	597	
			786	251. Melanetta, Boie..	804	598	
	Erimaturinae	811	252. Pelionetta, Kaup..	804	600	
			811	253. Oidemia, Flem....	804	601	
			811	254. Somateria, Leach	808	602	
			811	255. Erimatura, Bon..	811	604	
			812	256. Mergus, Linn....	813	606	
			812	257. Lophodytes, Reich	815	609	
			812	258. Mergellus, Selby..	817	611	
			812	613	

Sub-order B. GAVIAE. (Page 818.)

Tribe 1. TOTIPALMI. (Page 819.)

47. Pelecanidae....	868	259. Pelecanus, L.....	868	Cyrtopelicanus, Rch..	868	1	615	
48. Sulidae.....	871	260. Sula, Br.....	870	Oncorotulus, Wagl..	870	1	616	
					Sula, Br.....	871	1	617	
49. Tachypetidae...	873	261. Tachypetes, Vieill.	873	Dysporus, Ill.....	872	1	618	
					619	
50. Phalacrocoraci- dae.....	875	262. Graculus, L.....	875	Phalacrocorax, Br..	876	1	2	620
					Graculus, Bon.....	877	3	623	
					Urile, Bon.....	880	2	626	
51. Plotidae.....	883	Plotinae.....	883	263. Plotus, L.....	883	1	628	
52. Phaetonidae...	885	Phaetoninae.....	885	264. Phaeton, Linn....	885	1	629	

II. LIST OF SPECIES.¹

	Page.
1. <i>Cathartes aura</i> , (Linn.) Illig. Turkey Buzzard. All of North America, except the Arctic regions	4
2. <i>Cathartes californianus</i> , (Shaw,) Cuvier. California Vulture. Western North America	5
3. <i>Cathartes atratus</i> , (Bartram,) Less. Black Vulture. Southern North America. Central America to Chile	5
(4.) <i>Cathartes burrovianus</i> , Cassin. Burroughs' Vulture. Mexico, Vera Cruz, Mazatlan.	6
5. <i>Falco (Falco) anatum</i> , Bonaparte. Duck Hawk. North America, east of the Rocky mountains	7
6. <i>Falco (Falco) nigriceps</i> , Cassin. Western North and South America	8
7. <i>Falco (Hypotriorchis) columbarius</i> , Linn. Pigeon Hawk. Temperate North America, Mexico, Central America, Northern South America	9
8. <i>Falco (Hypotriorchis) aurantius</i> , Gmelin. Mexico, South America	10
9. <i>Falco (Hypotriorchis) femoralis</i> , Temminck. New Mexico, Mexico, South America	11
10. <i>Falco (Gennaia) polyagrus</i> , Cassin. Prairie Falcon. Western North America	12
11. <i>Falco (Hierofalco) candicans</i> , Gmelin. Jer Falcon. Northern North America; Greenland	13
12. <i>Falco (Hierofalco) islandicus</i> , Sabine. Jer Falcon. Northern North America; Greenland	13
13. <i>Falco (Tinnunculus) sparverius</i> , Linn. Sparrow Hawk. Entire continent of America	13
14. <i>Astur atricapillus</i> , (Wils.) Bonap. Goshawk. North America; chiefly in the north-western portions	15
15. <i>Accipiter cooperii</i> , Bonap. Cooper's Hawk. All of temperate North America	16
16. <i>Accipiter mexicanus</i> , Swains. Western N. America	17
17. <i>Accipiter fuscus</i> , (Gmelin,) Bon. Sharp-shinned Hawk. Throughout North America and Mexico	18
18. <i>Buteo swainsoni</i> , Bonap. Swainson's Buzzard. Northern and Western N. America	19
19. <i>Buteo bairdii</i> , Hoy. Baird's Buzzard. Northern and Western N. America	21
20. <i>Buteo calurus</i> , Cassin. Red-tailed Black Hawk. Western N. America	22
21. <i>Buteo (Leucopternis) insignatus</i> , Cassin. Western N. America	23
22. <i>Buteo (Leucopternis) harlani</i> , (Aud.) Bon. Harlan's Buzzard. Western N. America	24
23. <i>Buteo (Poecilopternis) borealis</i> , (Gmelin,) Vieill. Red-tailed Hawk. Eastern North America; fur countries; Cuba, Jamaica	25
24. <i>Buteo (Poecilopternis) montanus</i> , Nuttall. Western red-tailed Hawk. Western N. America	26
25. <i>Buteo (Poecilopternis) lineatus</i> , (Gmelin,) Jardine. Red-shouldered Hawk. Eastern and Northern N. America	28

¹When authorities are inclosed in parentheses it shows that the species was first described under a different genus. A second authority (or a single one uninclosed) is that of the name as adopted. Extra limital species have their current number similarly inclosed

	Page.
26. <i>Buteo (Poecilopternis) elegans</i> , Cassin. Western North America	28
27. <i>Buteo (Poecilopternis) pennsylvanicus</i> , (Wilson,) Bonap. Broad-winged Hawk. Eastern North America	29
28. <i>Buteo (Poecilopternis) oxypterus</i> , Cassin. New Mexico	30
29. <i>Buteo (Tachytriorchis) cooperi</i> , Cassin. California	31
30. <i>Archibuteo lagopus</i> , (Brünnich,) Gray. Rough-legged Hawk. All of temperate North America and Europe	32
31. <i>Archibuteo sancti-johannis</i> (Gmelin,) Gray. Black Hawk. Eastern and Northern North America	33
32. <i>Archibuteo ferrugineus</i> , (Licht.) Gray. California Squirrel Hawk. Western North America	34
33. <i>Asturina nitida</i> , (Lath.) Bonap. Northern Mexico and South America	35
34. <i>Nauclerus furcatus</i> , (Linn.) Vigors. Swallow-tailed Hawk. Eastern United States to the Mississippi, northward to Pennsylvania and Wisconsin; accidental in Europe	36
35. <i>Elanus leucurus</i> , (Vieillot,) Bonap. White-tailed Hawk. Southern and Western States and South America	37
36. <i>Ictinia mississippiensis</i> , (Wilson,) Gray. Mississippi Kite. Southern States, Texas, and New Mexico	37
37. <i>Rostrhamus sociabilis</i> , (Vieillot,) D'Orbigny. Black Kite. Florida and southward	38
38. <i>Circus hudsonius</i> , (Linn.) Vieillot. Marsh Hawk. All of North America and Cuba	38
39. <i>Aquila canadensis</i> , (Linn.) Cassin. Golden Eagle; Ring-tailed Eagle. All of North America	41
40. <i>Haliaeetus pelagicus</i> , (Pallas,) Siebold. Northern Sea Eagle. Russian American Islands; Japan	42
41. <i>Haliaeetus washingtonii</i> , (Aud.) Jard. Washington Eagle. Kentucky	42
42. <i>Haliaeetus albicilla</i> , (Linn.) Cuv. Gray Sea Eagle. Greenland; Europe	43
43. <i>Haliaeetus leucocephalus</i> , (Linn.) Savigny. Bald Eagle. All of temperate North America. Accidental in Europe	43
44. <i>Pandion carolinensis</i> , (Gmelin,) Bon. Fish Hawk. Throughout temperate North America	44
45. <i>Polyborus tharus</i> , (Molina,) Cassin. Caracara Eagle. Southern North America, Florida, Texas, Mexico	45
46. <i>Craxirex unicinctus</i> , (Temm.) Cassin. Harris' Buzzard. Southern States, Mexico, South America	46
47. <i>Strix pratincola</i> , Bonap. Barn Owl. Throughout temperate North America	47
48. <i>Bubo virginianus</i> , (Gmelin,) Bon. Great Horned Owl. The whole of North America. Runs into varieties, <i>atlanticus</i> , <i>pacificus</i> , <i>arcticus</i> , <i>magellanicus</i>	49
49. <i>Scops asio</i> , (Linn.) Bonap. Mottled Owl. The whole of temperate America, Greenland	51
50. <i>Scops m'callii</i> , Cassin. Western Mottled Owl. Western and southwestern North America	52
51. <i>Otus wilsonianus</i> , Lesson. Long-Eared Owl. The whole of temperate North America	53
52. <i>Brachyotus cassinii</i> , Brewer. Short-Eared Owl. The whole of temperate North America; Greenland; Cuba	54

73. 11710. 8. 21a - 2169. 8. 2a -

74. 110. 9. 1a.

75. 8087. 8. 7. 3. 8078. 9. 2. 2. 1.

76. 1291. 8. 1a.

77. 1291. 9. 1a.

LIST OF SPECIES.

XVII

	Page.
53. <i>Syrnium cinereum</i> , (Gmelin,) Aud. Great Gray Owl. Northern North America	56
54. <i>Syrnium nebulosum</i> , (Forster,) Gray. Barred Owl. Eastern North America; Fort Tejon, Cal.	56
55. <i>Nyctale richardsoni</i> , Bonap. Sparrow Owl. Northern North America; Canada	57
56. <i>Nyctale albifrons</i> , (Shaw,) Cassin. Kirtland's Owl. Northern North America; Wisconsin; Canada	57
57. <i>Nyctale acadica</i> , (Gmelin,) Bon. Saw-whet Owl. The whole of temperate North America; Fort Tejon, Cal.	58
58. <i>Athene hypugaea</i> , Bonap. Burrowing Owl. From Mississippi river to the Rocky Mountains	59
59. <i>Athene cunicularia</i> , (Molina,) Bon. Burrowing Owl. North America, west of the Rocky Mountains; South America	60
60. <i>Glaucidium gnoma</i> , (Wagler,) Cassin. Pigmy Owl. Oregon; California; Mexico	62
61. <i>Nyctea nivea</i> , (Daud.) Gray. Snowy Owl. Northern regions of both continents; Canada; Greenland; South Carolina; Bermuda	63
62. <i>Surnia ulula</i> , (Linn.) Bonap. Hawk Owl. Northern regions of both continents; Canada	64
63. <i>Conurus carolinensis</i> , (Linn.) Kuhl. Parakeet. Southern and southwestern States, as far west as the Missouri	67
(64.) <i>Rhynchopsitta pachyrhyncha</i> , (Sw.) Bon. Thick-billed Parrot. Rio Grande, Texas??	66
65. <i>Trogon mexicanus</i> , Swainson. Mexican Trogon. Northeastern Mexico to Rio Grande.	69
66. <i>Crotophaga rugirostris</i> , Sw. Black Parrot. Florida to Brazil	71
67. <i>Crotophaga ani</i> , Linn. Ani. Maritime parts of the southeastern United States and south to Brazil	72
68. <i>Geococcyx californianus</i> , (Less.) Baird. Chaparral Cock. Middle Texas, New Mexico, and California, to Central Mexico	73
69. <i>Coccyzus americanus</i> , (Linn.) Bonap. Yellow-billed Cuckoo. Eastern United States to the Missouri plains	76
70. <i>Coccyzus erythrophthalmus</i> , (Wils.) Bon. Black-billed Cuckoo. United States to the Missouri plains	77
71. <i>Coccyzus minor</i> , (Gmelin,) Cab. Mangrove Cuckoo. Florida keys to West Indies	78
72. <i>Campephilus principalis</i> , (Linn.) Gray. Ivory-billed Woodpecker. Southern Atlantic and Gulf States.	81
(73.) <i>Campephilus imperialis</i> , (Gould,) Gray. Imperial Woodpecker. Chiefly Central America and southwestern Mexico	82
74. <i>Picus (Trichopicus) villosus</i> , Linn. Hairy Woodpecker. Northern and western regions; var. <i>major</i> , <i>medius</i> , <i>minor</i>	84
75. <i>Picus (Trichopicus) harrisi</i> , Aud. Harris' Woodpecker. From the Pacific coast to the eastern slope of the Rocky Mountains	87
76. <i>Picus (Trichopicus) pubescens</i> , Linn. Downy Woodpecker. Eastern United States, towards the eastern slope of the Rocky Mountains	89
77. <i>Picus (Trichopicus) gairdneri</i> , Aud. Gairdner's Woodpecker. From Pacific coast to eastern base of Rocky Mountains	91

	Page.
78. <i>Picus (Dyctiopicus) nuttalli</i> , Gambel. Nuttall's Woodpecker. Coast region of California	93
79. <i>Picus (Dyctiopicus) scalaris</i> , Wagler. Rocky mountains and its slope, south of 35th parallel, to Yucatan	94
80. <i>Picus (Phrenopicus) borealis</i> , Vieill. Red-cockaded Woodpecker. Southern States . .	96
81. <i>Picus (Xenopicus) albolarvatus</i> , (Cass.) Baird. White-headed Woodpecker. Cascade mountains of Oregon, and southward into California	96
82. <i>Picoides arcticus</i> , (Sw.) Gray. Black-backed three-toed Woodpecker. Northern portions of the United States to the Arctic regions, from the Atlantic to the Pacific	98
83. <i>Picoides hirsutus</i> , (Vieill.) Gray. Banded three-toed Woodpecker. Arctic regions of North America	98 6
84. <i>Picoides dorsalis</i> , Baird. Striped three-toed Woodpecker. Laramie peak, Rocky Mountains	100
85. <i>Sphyrapicus varius</i> , (Linn.) Baird. Yellow-bellied Woodpecker. Atlantic ocean to the eastern slopes of Rocky Mountains, Greenland	103
86. <i>Sphyrapicus nuchalis</i> , Baird. Red-throated Woodpecker. Rocky mountains	103, 921
87. <i>Sphyrapicus ruber</i> , (Gm.) Baird. Red-breasted Woodpecker. Pacific slope of the United States	104
88. <i>Sphyrapicus williamsonii</i> , (Newb.) Baird. Williamson's Woodpecker. Rocky Mountains to the Cascade mountains	105
89. <i>Sphyrapicus thyroideus</i> , (Cass.) Baird. Brown-headed Woodpecker. Cascade and coast range of California and Oregon	106
90. <i>Hylatomus pileatus</i> , (Linn.) Baird. Black Woodcock. North America, from Atlantic to Pacific	107 5
91. <i>Centurus carolinus</i> , (Linn.) Bon. Red-bellied Woodpecker. North America, from Atlantic coast to the eastern slope of the Rocky Mountains	109
92. <i>Centurus flaviventris</i> , Sw. Yellow-bellied Woodpecker. Rio Grande region of the United States, south into Mexico	110
93. <i>Centurus uropygialis</i> , Baird. Gila Woodpecker. Lower Colorado river of the west . .	111
94. <i>Melanerpes erythrocephalus</i> , (Linn.) Sw. Red-headed Woodpecker. North America, from the Atlantic to the eastern slope of the Rocky Mountains . .	113
95. <i>Melanerpes formicivorus</i> , (Sw.) Bon. California Woodpecker. Coast region of California, and south; in northern Mexico, eastward almost to the Gulf of Mexico; also on upper Rio Grande	114
96. <i>Melanerpes torquatus</i> , (Wils.) Bonap. Lewis' Woodpecker. Western America, from Black Hills to Pacific	115
97. <i>Colaptes auratus</i> , (Linn.) Sw. Flicker. Eastern North America to the eastern slopes of the Rocky Mountains; Greenland	118
98. <i>Colaptes mexicanus</i> , Swain. Red-shafted Flicker. Western North America, from the Black Hills to Pacific	120
98a. <i>Colaptes hybridus</i> , Baird. Cross between two preceding. Upper Missouri	122
99. ? <i>Colaptes chrysoides</i> , (Malher.) Baird. New Mexico	125
100. <i>Lampornis mango</i> , (Linn.) Sw. Black-throated Humming Bird. Florida?	130

Melanerpes angustirostris - - - - -

101

102

103

104

105

106

107 1022 ♂ Cal. 4. - 1021 ♀ Cal. 4. 6017 ♀ 1131 ♀ - 6018 ♂ 2m.

108 11570 2m. 11571 ♀ 2m. 11572 ♂ 2m.

109 8217 ♂ Mo. - 8455 ♀ - 1010 ♀ -

110 12955 ♀ 2m. 3241 ♀ 2m. 12958 ♂ 2m.

111 17100 ♀ - ♂. 8498. ♀ - ♂. 1016 ♀ ♀ -

112 6963 ♀ Mo. 2144 ♂ Mo.

113 21244 ♂ Cal. 5491 ♂ Cal. 12204 ♀ 21459 ♂ Cal. 6250 ♀ 6002 ♀ - 1117 ♂ 2m. 2134 ♂ Cal. 1222 ♂ 2m.

114

115

116

117

118

119

120

121

122

123

124

125

	Page.
101. <i>Trochilus colubris</i> , Linn. Ruby-throated Humming Bird. Eastern North America to the High Central Plains; south to Brazil	131
102. <i>Trochilus alexandri</i> , Bourc. and Mulsant. Black-chinned Humming Bird. Coast of California, southward	133
103. <i>Selasphorus rufus</i> , (Gm.) Sw. Red-backed Humming Bird. West coast of North America, and across from Gulf of California to the Upper Rio Grande Valley	134
104. <i>Selasphorus platycercus</i> , (Sw.) Gould. Broad-tailed Humming Bird. High lands of Mexico, through Rocky mountains to Fort Bridger, Utah	135
105. <i>Atthis anna</i> , (Less.) Reich. Anna Humming Bird. Coast region of California	137
106. <i>Atthis costae</i> , (Bourc.) Reich. Southern California and Colorado Basin	138
107. <i>Panyptila melanoleuca</i> , Baird. White-throated Swift. Colorado Basin, New Mexico	141
108. <i>Nephocetes niger</i> , (Gm.) Baird. Northern Swift. Northwestern America (Simiahmoo bay, W. T.) to West India islands.	142
109. <i>Chaetura pelagica</i> , (Linn.) Steph. Chimney Swallow. Eastern United States to slopes of Rocky mountains.	144
110. <i>Chaetura vauxii</i> , (Townsend.) De Kay. Oregon Swift. Pacific coast, from Puget's Sound to California	145
111. <i>Antrostomus carolinensis</i> , (Gm.) Gould. Chuck-will's Widow. South Atlantic and Gulf States	147
112. <i>Antrostomus vociferus</i> , (Wils.) Bonap. Whip-poor-will. Eastern United States to the Plains	148
113. <i>Antrostomus nuttalli</i> , (Aud.) Cassin. Nuttall's Whip-poor-will. High Central Plains to the Pacific coast	149
114. <i>Chordeiles popetue</i> , (Vieill.) Baird. Night Hawk. North America generally.	151
115. <i>Chordeiles henryi</i> , Cassin. Western Night Hawk. Rocky mountains; north to Fort Bridger. Fort Laramie	153, 152
116. <i>Chordeiles texensis</i> , Lawrence. Texas Night-hawk. Rio Grande valley and south; west to Gulf of California	154
117. <i>Ceryle (Megaceryle) alcyon</i> , (Linn.) Boie. Belted King-fisher. Entire continent of North America	158
118. <i>Ceryle (Chloroceryle) americana</i> , (Gm.) Boie. Texas King-fisher. Rio Grande region of Texas and southward	159
119. <i>Momotus caerulescens</i> , Gould. Saw-bill. Mexico to Rio Grande valley	161
120. <i>Pachyrhynchus aglaiae</i> , Lafresnaye. Rose-throated Flycatcher. Mexico to Rio Grande	64
121. <i>Bathmidivus major</i> , Cab. Northeastern Mexico to Rio Grande?	166
122. <i>Milvulus tyrannus</i> , (Linn.) Bon. Fork-tailed Fly-catcher. South America to Vera Cruz; accidental in the United States	168
123. <i>Milvulus forficatus</i> , (Gm.) Sw. Scissor-tail. Central Texas to Mexico	169
124. <i>Tyrannus carolinensis</i> , (Linn.) Baird. King Bird. Eastern North America to Rocky mountains	171
125. <i>Tyrannus dominicensis</i> , (Briss.) Rich. Grey King Bird. South Carolina coast, accidental; Florida Keys and West Indies	172

	Page.
126. <i>Tyrannus verticalis</i> , Say. Arkansas Flycatcher. Western North America, from the High Central Plains to the Pacific	173
127. <i>Tyrannus vociferans</i> , Sw. Cassin's Flycatcher. Valley of Gila, eastward to Pecos river, Texas; and from South Pass into Mexico, on table-lands	174
128. <i>Tyrannus couchii</i> , Baird. Couch's Flycatcher. Northeastern Mexico to Rio Grande	175
(129.) <i>Tyrannus melancholicus</i> , Vieill. Southern Mexico, Central and South America	176
130. <i>Myiarchus crinitus</i> , (Linn.) Cab. Great Crested Flycatcher. Eastern North America to the Missouri and south to eastern Texas	178
131. <i>Myiarchus mexicanus</i> , (Kaup,) Baird. Ash-throated Flycatcher. Coast of California and across by Valley of Gila and Rio Grande to northeastern Mexico	179
(132.) <i>Myiarchus cooperi</i> , (Kaup,) Baird. Mexico	180
133. <i>Myiarchus lawrencii</i> , (Giraud,) Baird. Lawrence's Flycatcher. Northeastern Mexico to the Rio Grande	181
134. <i>Sayornis nigricans</i> , (Sw.) Bon. Black Flycatcher. California coast and across by Valley of Gila and upper Rio Grande to New Leon, and south	183
135. <i>Sayornis fuscus</i> , (Gm.) Baird. Pewee. Eastern North America	184
136. <i>Sayornis sayus</i> , (Bon.) Baird. Say's Flycatcher. Upper Missouri river and Central High Plains westward to the Pacific and south to Mexico	185
137. <i>Contopus borealis</i> , (Swainson,) Baird. Olive-sided Flycatcher. Rare on the Atlantic and Pacific coasts of the United States; Fort Bridger. Found in Greenland	188
138. <i>Contopus richardsonii</i> , (Sw.) Baird. Short-legged Pewee. High central dry plains to the Pacific; Rio Grande valley southward to Mexico; Labrador	189
139. <i>Contopus virens</i> , (Linn.) Cab. Wood Pewee. Eastern North America to the borders of the High Central Plains; south to New Granada	190
140. <i>Empidonax traillii</i> , (Aud.) Baird. Traill's Flycatcher. Eastern United States, and south to Mexico	193
141. <i>Empidonax pusillus</i> , (Sw.) Cab. High Central Plains to the Pacific; Fur countries; southward into Mexico	194
142. <i>Empidonax minimus</i> , Baird. Least Flycatcher. Eastern United States to Fort Bridger	195
143. <i>Empidonax acadicus</i> , (Gm.) Baird. Small Green-crested Flycatcher. Eastern United States to the Mississippi	197
144. <i>Empidonax flaviventris</i> , Baird. Yellow-bellied Flycatcher. Eastern United States generally; perhaps replaced on the Pacific by a closely allied species, <i>E. difficilis</i>	198
145. <i>Empidonax hammondi</i> , (de Vesey,) Baird. Vicinity of Fort Tejon to Los Angeles, Cal.	199
146. <i>Empidonax obscurus</i> , (Sw.) Baird. Rocky mountains to Fort Bridger, Utah; Fort Yuma	200
147. <i>Pyrocephalus rubineus</i> , (Bodd.) Gray. Red Flycatcher. Valleys of Rio Grande and Gila, southward	201
148. <i>Turdus (Turdus) mustelinus</i> , Gm. Wood Thrush. Eastern United States to Missouri river; south to Guatemala	212

	Page.
149. <i>Turdus (Turdus) pallasi</i> , Cab. Hermit Thrush. Eastern North America to the Mississippi river	212
var. <i>silens</i> , Swainson. High lands of Mexico to Fort Bridger	213
150. <i>Turdus (Turdus) nanus</i> , Aud. Dwarf Thrush. Pacific coast of North America to Rocky mountains	213
151. <i>Turdus (Turdus) fuscescens</i> , Stephens. Wilson's Thrush. Eastern North America to the Missouri; north to fur countries	214
152. <i>Turdus (Turdus) ustulatus</i> , Nuttall. Coast region of Oregon and Washington Territories	215
153. <i>Turdus (Turdus) swainsoni</i> , Cab. Olive-backed Thrush. Eastern North America to Fort Bridger; south to Mexico and Peru; north to Greenland; accidental in Europe and Siberia	216
154. <i>Turdus (Turdus) aliciae</i> , Baird. Gray-cheeked Thrush. Mississippi region to the Missouri	217
155. <i>Turdus (Planesticus) migratorius</i> , Linn. Robin. Continent of North America to Mexico	218
156. <i>Turdus (Ixoreus) naevius</i> , Bon. Varied Thrush. Pacific coast of North America; accidental on Long Island and near Boston	219
(157.) <i>Saxicola oenanthe</i> , (Linn.) Bechst. Stone Chat. Greenland; accidental in northern part of North America; common in Europe	220
158. <i>Sialia sialis</i> , (Linn.) Baird. Blue Bird. Eastern North America to west of Missouri river; Fort Laramie	222
159. <i>Sialia mexicana</i> , Sw. Western Blue Bird. Pacific coast of North America, and along Valley of Gila to upper Rio Grande, and south	223
160. <i>Sialia arctica</i> , Sw. Rocky mountain Blue Bird. Upper Missouri to Rocky mountains range, and south to Mexico; rare on the coast of California	224
161. <i>Regulus calendula</i> , (Linn.) Licht. Ruby-crowned Wren. United States, from Atlantic to Pacific.	226
162. <i>Regulus satrapa</i> , Licht. Golden-crested Wren. Northern parts of United States; from Atlantic to Pacific; on west coast only noticed on Puget's Sound	227
163. <i>Regulus cuvieri</i> , Aud. Cuvier's Golden Crest. Pennsylvania	228
164. <i>Hydrobata mexicana</i> , (Bon.) Baird. Water Ouzel. Rocky mountains and west from British America to Mexico. Fort Tejon	229
165. <i>Anthus ludovicianus</i> , (Gm.) Licht. Tit-lark. North America generally; Greenland; accidental in Europe	232
166. <i>Neocorys spragueii</i> , (Aud.) Sclater. Missouri Skylark. About Fort Union, Nebraska	234
167. <i>Mniotilta varia</i> , (Linn.) Vieill. Black and white Creeper. Eastern North America to Missouri river; south to Guatemala. Var. <i>longirostris</i> , Baird. Florida	235
168. <i>Parula americana</i> , (Linn.) Bonap. Blue yellow-backed Warbler. Eastern North America to Missouri river; south to Guatemala	238
169. <i>Protonotaria citrea</i> , (Bodd.) Baird. Prothonotary Warbler. South Atlantic and Gulf States to mouth of Ohio north, and the Missouri river west; West Indies	239

	Page.
170. <i>Geothlypis trichas</i> , (Linn.) Cab. Maryland Yellow-throat. North America from Atlantic to Pacific.....	241
(171.) <i>Geothlypis velatus</i> , (Vieill.) Cab. Green-headed Warbler. West Indies and South America	243
172. <i>Geothlypis philadelphia</i> , (Wils.) Baird. Mourning Warbler. Eastern North America as far west as Independence, Mo.....	243
173. <i>Geothlypis macgillivrayi</i> , (Aud.) Baird. Macgillivray's Warbler. Pacific coast of North America, south to Gulf of California, and across to Monterey, Mexico. In Rocky mountains to Fort Laramie.....	244
174. <i>Oporornis agilis</i> , (Wils.) Baird. Connecticut Warbler. Eastern United States	246
175. <i>Oporornis formosus</i> , (Wils.) Baird. Kentucky Warbler. Eastern United States as far west as Fort Riley; south to Guatemala	247
176. <i>Icteria viridis</i> , (Gm.) Bonap. Yellow-breasted Chat. Eastern United States to the Missouri; south to Guatemala	248
177. <i>Icteria longicauda</i> , Lawr. Long-tailed Chat. High Central Plains of United States to the Pacific; south into Mexico.....	249
178. <i>Helmintherus vermivorus</i> , (Gm.) Bonap. Worm-eating Warbler. Eastern United States to Missouri river; south to Guatemala.....	252
179. <i>Helmintherus swainsonii</i> , (Aud.) Sw. Swainson's Warbler. South Atlantic States....	252
180. <i>Helminthophaga pinus</i> , (Linn.) Baird. Blue-winged Yellow Warbler. Eastern United States to the Missouri; south to Guatemala.....	254
181. <i>Helminthophaga chrysoptera</i> , (Linn.) Baird. Golden-winged Warbler. Eastern United States to the Missouri; Bogota.....	255
182. <i>Helminthophaga bachmani</i> , (Aud.) Cab. Bachman's Warbler. South Atlantic States; Cuba.....	255
183. <i>Helminthophaga ruficapilla</i> , (Wils.) Baird. Nashville Warbler. Eastern North America to the Missouri; Fort Tejon, Cal.; Greenland.....	256
184. <i>Helminthophaga celata</i> , (Say.) Baird. Orange-crowned Warbler. Mississippi river to the Pacific; south to northern Mexico.....	257
185. <i>Helminthophaga peregrina</i> , (Wils.) Cab. Tennessee Warbler. Eastern United States to the Missouri.....	258
186. <i>Seiurus aurocapillus</i> , (Linn.) Sw. Golden-crowned Thrush. Eastern North America to the Missouri.....	260
187. <i>Seiurus noveboracensis</i> , (Gm.) Nutt. Water Thrush. Eastern United States to the Missouri, and south to Guatemala, perhaps to Brazil.....	261
188. <i>Seiurus ludovicianus</i> , (Vieill.) Bon. Large-billed Water Thrush. Eastern United States to the Missouri; south to Mexico.....	262
189. <i>Dendroica virens</i> , (Gm.) Baird. Black-throated Green Warbler. Eastern United States to the Missouri; south to Guatemala; Greenland.....	267
190. <i>Dendroica occidentalis</i> , (Townsend.) Baird. Western Warbler. Pacific coast.....	268
191. <i>Dendroica townsendii</i> , (Nutt.) Baird. Pacific coast, North America, to the Rocky Mountains; south to Mexico and Guatemala.....	269
192. <i>Dendroica nigrescens</i> , (Townsend.) Baird. Black-throated Gray Warbler. Pacific coast United States; Fort Thorn, New Mexico	270

	Page.
193. <i>Dendroica canadensis</i> , (Linn.) Baird. Black-throated Blue Warbler. Eastern United States to the Missouri; south to the West Indies	271
194. <i>Dendroica coronata</i> , (Linn.) Gray. Yellow Rump. Eastern North America to the Missouri plains; stragglers seen on Puget's Sound	272
195. <i>Dendroica audubonii</i> , (Townsend) Baird. Audubon's Warbler. Pacific coast of United States to Rocky mountains; south to Mexico	273
196. <i>Dendroica Blackburniae</i> , (Gm.) Baird. Blackburnian Warbler. Eastern North America to the Missouri; south to Guatemala	274
197. <i>Dendroica castanea</i> , (Wils.) Baird. Bay Breasted Warbler. Eastern United States to the Missouri; south to Guatemala	276
198. <i>Dendroica pinus</i> , (Wils.) Baird. Pine Creeping Warbler. Eastern United States to the Missouri	277
199. <i>Dendroica montana</i> , (Wils.) Baird. Blue Mountain Warbler. Blue mountains of Virginia	278
200. <i>Dendroica pennsylvanica</i> , (Linn.) Baird. Chestnut-sided Warbler. Eastern United States to the Missouri	279
201. <i>Dendroica caerulea</i> , (Wils.) Baird. Blue Warbler. Eastern United States to the Missouri river	280
202. <i>Dendroica striata</i> , (Forster.) Baird. Black Poll Warbler. Eastern North America to the Missouri high plains; Cuba; Greenland	280
203. <i>Dendroica aestiva</i> , (Gm.) Baird. Yellow Warbler. United States from Atlantic to Pacific; south to Guatemala and West Indies	282
204. <i>Dendroica maculosa</i> , (Gm.) Baird. Black and Yellow Warbler. Eastern United States to the Missouri; south to Guatemala	284
205. <i>Dendroica kirtlandii</i> , Baird. Kirtland's Warbler. Northern Ohio	286
206. <i>Dendroica tigrina</i> , (Gm.) Baird. Cape May Warbler. Eastern United States to the Mississippi; Cuba	286
207. <i>Dendroica carbonata</i> , (Aud.) Baird. Kentucky	287
208. <i>Dendroica palmarum</i> , (Gm.) Baird. Yellow Red Poll. Eastern United States to the Mississippi, and Red river of the north	288
209. <i>Dendroica superciliosa</i> , (Bodd.) Baird. Yellow-throated Warbler. Eastern United States as far as Pennsylvania and Ohio to the Missouri; south to Mexico	289
210. <i>Dendroica discolor</i> , (Vieill.) Baird. Prairie Warbler. Atlantic States, as far north as New York	290
211. <i>Mniotiltus mitratus</i> , (Gm.) Aud. Hooded Warbler. Eastern United States to the Missouri; south to Guatemala	292
212. <i>Mniotiltus minutus</i> , (Wils.) Baird. Small-headed Flycatcher. Eastern Atlantic States	293
213. <i>Mniotiltus pusillus</i> , (Wils.) Bon. Green Black-cap Flycatcher. United States from Atlantic to Pacific; south to Guatemala	293
214. <i>Mniotiltus canadensis</i> , (Linn.) Aud. Canada Flycatcher. Eastern United States to the Mississippi; south to Guatemala	294
215. <i>Mniotiltus bonapartei</i> , Aud. Bonaparte's Flycatcher. Louisiana	295
(216.) <i>Cardellina rubra</i> , (Sw.) Bon. Vermilion Flycatcher. Northern Mexico	296

	Page.
217. <i>Setophaga ruticilla</i> , (Linn.) Sw. Redstart. Eastern United States to Fort Bridger ; West Indies in winter	297
218. <i>Setophaga picta</i> , Sw. Northern Mexico ; Rio Grande valley	298
(219.) <i>Setophaga miniata</i> , Sw. Northern Mexico (Rio Grande valley) to Guatemala	299
220. <i>Pyrranga rubra</i> , (Linn.) Vieill. Scarlet Tanager. Eastern United States to the Mis- souri river	300
221. <i>Pyrranga aestiva</i> , (Linn.) Vieillot. Summer Red Bird. South Atlantic and Gulf States, through Texas, and south to Guatemala	301
222. <i>Pyrranga hepatica</i> , Sw. Rocky mountains of New Mexico, southward	302
223. <i>Pyrranga ludoviciana</i> , (Wils.) Bon. Louisiana Tanager. From the Black Hills to the Pacific ; south to Mexico	303
(224.) <i>Euphonia elegantissima</i> , (Bon.) Gray. Northern Mexico to Guatemala. Califor- nia?	304
225. <i>Hirundo horreorum</i> , Barton. Barn Swallow. North America from Atlantic to Pacific	308
226. <i>Hirundo lunifrons</i> , Say. Cliff Swallow. North America from Atlantic to Pacific	309
227. <i>Hirundo bicolor</i> , Vieill. White-bellied Swallow. North America from Atlantic to Pa- cific	310
228. <i>Hirundo thalassina</i> , Sw. Violet Green Swallow. Rocky mountains to Pacific. South Mexico ; east to Saltillo, Mexico	311
229. <i>Cotyle riparia</i> , (Linn.) Boie. Bank Swallow. North America generally	313
230. <i>Cotyle (Stelgidopteryx) serripennis</i> , (Aud.) Bon. Rough-winged Swallow. United States from Atlantic to Pacific	313
231. <i>Progne purpurea</i> , (Linn.) Boie. Purple Martin. North America generally	314
<i>Progne</i> ———? Indian Key, Florida	App. 923
232. <i>Ampelis garrulus</i> , Linn. Wax Wing. Northern parts of both continents. Seen in the United States only in severe winters, except along the great lakes. In the Mississippi valley south to Fort Riley. "Millions" seen north of Fort Bridger	317
233. <i>Ampelis cedrorum</i> , (Vieill.) Baird. Cedar Bird. North America generally ; south to Guatemala	318
234. <i>Phainopepla nitens</i> , (Swainson,) Sclater. Valley of Gila and southern Colorado to up- per Rio Grande ; west to Fort Tejon ; east to Coahuila, Mexico	320, 923
235. <i>Myiadestes townsendii</i> , (Aud.) Cab. Townsend's Flycatcher. United States, from Rocky Mountains and Black Hills to the Pacific ; south to the borders of Mexico	321
236. <i>Collyrio borealis</i> , (Bon.) Baird. Great Northern Shrike. Northern regions, from At- lantic to Pacific ; in winter south, through most of the United States	324
237. <i>Collyrio ludovicianus</i> , (Linn.) Baird. Loggerhead Shrike. South Atlantic and Gulf States	325
238. <i>Collyrio excubitoroides</i> , (Sw.) Baird. White-rumped Shrike. Missouri plains and fur countries to Pacific coast ; eastward into Wisconsin, Illinois, and Michigan (?)	327

	Page.
239. <i>Collyrio elegans</i> , (Sw.) White-winged Shrike. Western America	328
240. <i>Vireo (Vireosylva) olivaceus</i> , (Linn.) Vieill. Red-eyed Flycatcher. Eastern United States to Fort Bridger, Utah; in Texas to Devil's river; south to Guatemala; Greenland	321
241. <i>Vireo (Vireosylva) flavoviridis</i> , Cassin. Northern Mexico to Panama	332
242. <i>Vireo (Vireosylva) virescens</i> , Vieill. Bartram's Vireo. Central and eastern South America; Atlantic United States	333
243. <i>Vireo (Vireosylva) altiloquus</i> , (Vieill.) Gray. Whip Tom Kelly. Coast of southern Florida and the West Indies	334
244. <i>Vireo philadelphicus</i> , Cassin. Pennsylvania to Wisconsin	335
245. <i>Vireo gilvus</i> , (Vieill.) Bon. Warbling Flycatcher. Atlantic to Pacific coast of United States; var. <i>swainsonii</i> ; Columbia river	335
246. <i>Vireo belli</i> , Aud. Bell's Vireo. Missouri and eastern Texas	337
247. <i>Vireo atricapillus</i> , Woodh. Black-headed Flycatcher. Devil's river, Texas	337
248. <i>Vireo (Lanivireo) noveboracensis</i> , (Gm.) Bon. White-eyed Vireo. Eastern United States to the Missouri and throughout Texas	338
249. <i>Vireo (Lanivireo) huttoni</i> , Cass. Hutton's Flycatcher. South California, across by valley of Gila to northeastern Mexico	339
250. <i>Vireo (Lanivireo) solitarius</i> , (Wils.) Vieill. Blue-headed Flycatcher. United States from Atlantic to the Pacific	340
251. <i>Vireo (Lanivireo) cassinii</i> , De Vesev. Cassin's Vireo. Fort Tejon, California	340
252. <i>Vireo (Lanivireo) flavifrons</i> , Vieill. Yellow-throated Flycatcher. Eastern United States to Missouri; south to Central America	341
253. <i>Mimus polyglottus</i> , (Linn.) Boie. Mocking Bird. Southern United States from Atlantic to High Central Plains. Perhaps replaced by another species to the Pacific. (<i>M. caudatus</i>)	344
254. <i>Mimus carolinensis</i> , (Linn.) Gray. Cat Bird. Eastern United States to Fort Bridger	346
255. <i>Oreoscoptes montanus</i> , (Townsend) Baird. Mountain Mocking Bird. Rocky mountains from Fort Bridger south to Mexico; along valley of Gila and Colorado; San Diego, California	347
256. <i>Harporhynchus redivivus</i> , (Gambel,) Cab. Coast of California	349
257. <i>Harporhynchus lecontei</i> , (Lawr.) Bonap. Fort Yuma, California	350
258. <i>Harporhynchus crissalis</i> , Henry. Fort Thorn to Fort Yuma	351
259. <i>Harporhynchus curvirostris</i> , (Sw.) Cab. Lower Rio Grande	351
259. ? <i>Harporhynchus vetula</i> , (Wagl.) New Mexico	352
260. <i>Harporhynchus longirostris</i> , (Lafres.) Cab. Lower Rio Grande, south through eastern Mexico	352
261. <i>Harporhynchus rufus</i> , (Linn.) Cab. Brown Thrush. Eastern North America to Missouri river, and perhaps to High Central Plains, unless replaced by a long-tailed variety, (<i>H. longicauda</i>)	353
262. <i>Campylorhynchus brunneicapillus</i> , (Lafres.) Gray. Valleys of Rio Grande and Gila, southward	355

	Page.
263. <i>Catherpes mexicanus</i> , (Sw.) Baird. White-throated Wren. Valley of Rio Grande, Colorado, and Gila, (but not on the coast of California,) south into Mexico	356
264. <i>Salpinctes obsoletus</i> , (Say.) Cab. Rock Wren. High Central Plains through the Rocky mountains to the Cascade range, (but not on the Pacific coast?) Fort Tejon	357
265. <i>Thriothorus ludovicianus</i> , (Gm.) Bon. Great Carolina Wren. Eastern United States to the Missouri; north to Pennsylvania; in Texas, to upper Rio Grande	361
266. <i>Thriothorus berlandieri</i> , Couch. Northeastern Mexico, towards the Rio Grande	362
267. <i>Thriothorus bewickii</i> , (Aud.) Bon. Bewick's Wren. North America, from Atlantic to Pacific; south to Mexico. The western bird possibly distinct, (<i>T. spilurus</i>)	363
268. <i>Cistothorus (Telmatoodytes) palustris</i> , (Wils.) Cab. Long-billed Marsh Wren. North America, from Atlantic to Pacific; north to Greenland	364
269. <i>Cistothorus (Cistothorus) stellaris</i> , (Licht.) Cab. Short-billed Marsh Wren. Eastern United States to the Loup Fork of Platte	365
270. <i>Troglodytes aedon</i> , Vieill. House Wren. Eastern United States to the Missouri, or to the High Central Plains	367
271. <i>Troglodytes parkmani</i> , Aud. Parkman's Wren. Western America, from the High Central Plains and upper Missouri to the Pacific	367
272. <i>Troglodytes americanus</i> , Aud. Wood Wren. Eastern United States	368
273. <i>Troglodytes (Anorhwa) hyemalis</i> , (Wils.) Vieill. Winter Wren. North America generally	369
274. <i>Chamaca fasciata</i> , Gambel. Coast of California	370
275. <i>Certhia americana</i> , Bonap. American Creeper. Eastern North America	372
276. <i>Certhia mexicana</i> , Gloger. Rocky mountains to Pacific; Mexico	373
277. <i>Sitta carolinensis</i> , Gm. White-bellied Nuthatch. Eastern North America to the High Central Plains. West of this replaced by <i>S. aculeata</i>	374
278. <i>Sitta aculeata</i> , Cassin. Slender-bill Nuthatch. Pacific coast, and east towards the Rocky mountains	375
279. <i>Sitta canadensis</i> , Linn. Red-bellied Nuthatch. North America to the Rocky mountains; probably also to the Pacific	376
280. <i>Sitta pusilla</i> , Latham. Brown-headed Nuthatch. South Atlantic (and Gulf?) States	377
281. <i>Sitta pygmaea</i> , Vigors. California Nuthatch. Pacific coast and towards Rocky mountains; New Mexico	378
282. <i>Poliophtila caerulea</i> , (Linn.) Sclat. Blue-Gray Flycatcher. United States, from Atlantic to Missouri, and on the southern border from the Gulf of Mexico to the coast mountains of California, south to Guatemala	380
283. <i>Poliophtila plumbea</i> , Baird. Valley of Colorado and Gila	382
284. <i>Poliophtila melanura</i> , Law. Valley of Rio Grande and Gila; west to San Diego	382
285. <i>Lophophanes bicolor</i> , (Linn.) Bon. Tufted Titmouse. Eastern North America to the Missouri river, or else replaced there by a <i>L. missouriensis</i>	384

	Page.
286. <i>Lophophanes atricristatus</i> , Cassin. Black-crested Tit. Valley of Rio Grande and south	385
287. <i>Lophophanes inornatus</i> , (Gamb.) Cassin. Coast of California and southern Rocky mountains	386
288. <i>Lophophanes wollweberi</i> , Bon. Southern Rocky mountains, and south into Mexico, on the table-lands	386
289. <i>Parus septentrionalis</i> , Harris. Long-tailed Chickadee. Missouri river to the Rocky mountains, or else replaced there by <i>P. albescens</i>	389
290. <i>Parus atricapillus</i> , Linn. Black-cap Titmouse. Eastern North America, along the Atlantic border	390
291. <i>Parus occidentalis</i> , Baird. Western Titmouse. North Pacific coast of United States ..	391
(292.) <i>Parus meridionalis</i> , Sclater. Mexican Titmouse. Eastern Mexico	392
293. <i>Parus carolinensis</i> , Aud. Carolina Titmouse. South Atlantic States to Washington ..	392
294. <i>Parus montanus</i> , Gambel. Pacific coast of United States to the Rocky mountains ..	394
295. <i>Parus rufescens</i> , Towns. Chestnut-backed Tit. Pacific coast of the United States ..	394
296. <i>Parus hudsonicus</i> , Forster. Northeastern portions of North America to the North Atlantic States	395
(297.) <i>Psaltriparus melanotus</i> , (Hart.) Bon. Black-cheeked Tit. Eastern Mexico to the Rio Grande	396
298. <i>Psaltriparus minimus</i> , (Towns.) Bon. Least Tit. Pacific coast of United States	397
299. <i>Psaltriparus plumbeus</i> , Baird. Southern Rocky mountains	398
300. <i>Paroides flaviceps</i> , (Sund.) Baird. Rio Grande valley of Texas and Mexico. Fort Yuma	400
301. <i>Certhiola flaveola</i> , (Linn.) Sund. Yellow-rumped Creeper. Indian Key, Florida, and West Indies	924
302. <i>Eremophila cornuta</i> , (Wils.) Boie. Sky Lark. Everywhere on the prairies and desert plains of North America; Atlantic States in winter. A smaller variety on the western plains	403
303. <i>Hesperiphona vespertina</i> , (Cooper,) Bon. Evening Grosbeak. Pacific coast to Rocky mountains; Northern America east to Lake Superior	400
304. <i>Pinicola canadensis</i> , (Briss.) Cab. Pine Grosbeak. Arctic America, south to United States in severe winters	410
305. <i>Carpodacus purpureus</i> , (Gm.) Gray. Purple Finch. North America from Atlantic to the High Central Plains	412
306. <i>Carpodacus californicus</i> , Baird. Western Purple Finch. Pacific coast of United States	413
307. <i>Carpodacus cassinii</i> , Baird. Cassin's Purple Finch. Rocky mountains and valley of the Colorado. Fort Tejon?	414
308. <i>Carpodacus frontalis</i> , (Say,) Gray. House Finch. Rocky mountains to the Pacific. Perhaps several varieties	415
(309.) <i>Carpodacus haemorrhous</i> , Wagler. High lands of Mexico	417
310. <i>Chrysomitris magellanicus</i> , (Vieill.) Bon. Black-headed Gold-finch. South America. Accidental in United States, (Kentucky)	419
311. <i>Chrysomitris stanleyi</i> , (Aud.) Bon. Stanley's Gold-finch. California (?) Probably western Mexico	420

	Page.
312. <i>Chrysomitris yarrelli</i> , (Aud.) Bon. Yarrell's Gold-finch. California (?) Probably western Mexico	421
313. <i>Chrysomitris tristis</i> , (Linn.) Bon. Yellow Bird. North America generally	421
314. <i>Chrysomitris psaltria</i> , (Say.) Bon. Arkansas Finch. Southern Rocky mountains to the coast of California	422
315. <i>Chrysomitris mexicana</i> , (Sw.) Bon. Mexican Goldfinch. Mexican side of the valley of the Rio Grande, southward; Copper Mines of the Gila	423
316. <i>Chrysomitris lawrencii</i> , (Cassin.) Bon. Coast of California	424
317. <i>Chrysomitris pinus</i> , (Wils.) Bon. Pine Finch. North America, from Atlantic to Pacific	425
318. <i>Curvirostra americana</i> , Wils. Red Crossbill. North America generally, coming southward in winter. Resident in the mountains of Pennsylvania. Perhaps var. <i>mexicana</i> in Rocky mountains	426
319. <i>Curvirostra leucoptera</i> , (Gm.) Wils. White-winged Crossbill. Northern parts of North America generally	427
320. <i>Aegiothus linaria</i> , (Linn.) Cab. Lesser Red Poll. Throughout eastern North America, coming south in winter; Washington Territory	428
321. <i>Aegiothus canescens</i> , (Gould.) Cab. Mealy Red Poll. Greenland	429
322. <i>Leucosticte tephrocotis</i> , Sw. Gray-crowned Finch. Northern Rocky mountains; vicinity of Salt Lake City in winter	430
323. <i>Leucosticte griseinucha</i> , (Brand.) Bonap. Russian America, Aleutians	430
324. <i>Leucosticte arctous</i> , (Pall.) Bonap. Kamtschatka; Russian America	430
325. <i>Plectrophanes (Plectrophanes) nivalis</i> , (Linn.) Meyer. Snow Bunting. North America, from Atlantic to Pacific; south into the United States in winter	432
326. <i>Plectrophanes (Centrophanes) lapponicus</i> , (Linn.) Selby. Lapland Longspur. Eastern North America into United States in winter. Not yet found much west of the Missouri. Fort Laramie?	433
327. <i>Plectrophanes (Centrophanes) pictus</i> , Sw. Smith's Bunting. Prairies of Illinois in winter; in summer north to the Saskatchewan	434
328. <i>Plectrophanes (Centrophanes) ornatius</i> , Towns. Chestnut-collared Bunting. Plains of the Upper Missouri	435
329. <i>Plectrophanes (Centrophanes) melanomus</i> , Baird. Eastern slope of the Rocky mountains; Mexico, on the table-lands	436
330. <i>Plectrophanes (Rhynchophanes) maccownii</i> , Lawr. Eastern slopes of Rocky mountains; from Fort Thorn, N. M., as far east as the Black Hills, north of the Platte	437
331. <i>Centronyx bairdii</i> , (Aud.) Baird. Mouth of the Yellowstone	441
332. <i>Passerculus savanna</i> , (Wils.) Bon. Savannah Sparrow. Eastern North America to the Missouri plains	442
333. <i>Passerculus sandwichensis</i> , (Gm.) Baird. Northwestern coast, from the Columbia river to Russian America	444
334. <i>Passerculus anthinus</i> , Bon. Coast of California, near San Francisco; Russian America, Kodiak	445

	Page.
335. <i>Passerculus alaudinus</i> , Bon. Coast of California and lower Rio Grande of Texas and Mexico; Fort Bridger	446
336. <i>Passerculus rostratus</i> , (Cass.) Baird. Coast of California near San Diego.	446
337. <i>Poocetes gramineus</i> , (Gm.) Baird. Grass Finch. United States from Atlantic to the Pacific; or else one species to the high central plains, and another from this to the Pacific	447
338. <i>Coturniculus passerinus</i> , (Wils.) Bon. Yellow-winged Sparrow. Eastern United States to the high central plains (Loup Fork); also along the valley of Gila and Colorado	450
339. <i>Coturniculus henslowi</i> , (Aud.) Bon. Henslow's Bunting. Eastern United States as far north as Washington; westward to the Loup Fork of Platte	451
340. <i>Coturniculus lecontei</i> , (Aud.) Bon. Leconte's Bunting. Mouth of Yellowstone	452
341. <i>Ammodromus cauducutus</i> , (Gm.) Sw. Sharp-tailed Finch. Atlantic coast of the United States	453
342. <i>Ammodromus maritimus</i> , (Wils.) Sw. Sea-side Finch. Atlantic coast as far, at least, as Long Island	454
343. <i>Ammodromus samuelis</i> , Baird. California	455
344. <i>Chondestes grammaca</i> , (Say,) Bon. Lark Finch. From Wisconsin and the prairies of Michigan to Pacific coast; south to Texas and Mexico, on the plains	456
345. <i>Zonotrichia leucophrys</i> , (Forster,) Sw. White-crowned Sparrow. United States from Atlantic to the Rocky mountains, where they become mixed up with <i>Z. gambelii</i> ; Greenland.	458
346. <i>Zonotrichia gambelii</i> , (Nutt.) Gambel. Rocky mountains to the Pacific coast; Fort Bridger	460
347. <i>Zonotrichia coronata</i> , (Pallas.) Baird. Golden-crowned Sparrow. Pacific coast from Russian America to southern California; Black Hills of Rocky mountains?	461
348. <i>Zonotrichia querula</i> , (Nutt.) Gamb. Harris's Finch. Missouri river, above Fort Leavenworth	462
349. <i>Zonotrichia albicollis</i> , (Gm.) Bon. White-throated Sparrow. Eastern United States to the Missouri	463
(350.) <i>Junco cinereus</i> (Sw.) Cab. Mexico	465
351. <i>Junco dorsalis</i> , Henry. Fort Thorn, New Mexico	466
352. <i>Junco oregonus</i> , (Townsend.) Sclat. Oregon Snow Bird. Pacific coast of the United States to the eastern side of the Rocky mountains; stragglers as far east as Fort Leavenworth in winter, and Great Bend of Missouri	467
353. <i>Junco caniceps</i> , (Woodh.) Baird. Rocky mountains, from Black Hills to Fort Bridger and San Francisco; mountains of New Mexico	468
354. <i>Junco hyemalis</i> , (Linn.) Sclat. Snow Bird. Eastern United States to the Missouri, and as far west as Black Hills	468
355. <i>Poospiza bilineata</i> , (Cass.) Sclat. Black-throated Sparrow. Valley of Rio Grande and of Gila. (Big cañon of Colorado.)	470

356. <i>Poospiza belli</i> , (Cass.) Sclat. Bell's Finch. Southern California and valley of Gila and Colorado to Fort Thorn, and north to Fort Bridger, where very abundant	470
357. <i>Spizella monticola</i> , (Gm.) Baird. Tree Sparrow. Eastern North America to the Missouri; also on Pole creek and Little Colorado river, New Mexico	472
358. <i>Spizella pusilla</i> , (Wils.) Bon. Field Sparrow. Eastern North America to the Missouri river	473
359. <i>Spizella socialis</i> , (Wils.) Bon. Chipping Sparrow. North America from Atlantic to Pacific	473
360. <i>Spizella pallida</i> , (Sw.) Bon. Clay-colored Bunting. Upper Missouri river and High Central Plains to the Saskatchewan country	474
361. <i>Spizella breweri</i> , (Aud.) Cass. Brewer's Sparrow. Rocky mountains of United States to the Pacific coast	475
362. <i>Spizella atrigularis</i> , (Cab.) Baird. Black-chinned Sparrow. Mexico just south of the Rio Grande	476
363. <i>Melospiza (Melospiza) melodia</i> , (Wils.) Baird. Song Sparrow. Eastern United States to the High Central Plains.	477
364. <i>Melospiza (Melospiza) heermanni</i> , Baird. Heermann's Song Sparrow. Tejon Pass, California	478
365. <i>Melospiza (Melospiza) gouldii</i> , Baird. California.	479
366. <i>Melospiza (Melospiza) rufina</i> , (Brandt.) Baird. Pacific coast of the United States to Russian America.	480
367. <i>Melospiza (Melospiza) fallax</i> , Baird. Rocky mountain region from Fort Thorn to the Colorado	481
368. <i>Melospiza (Melospiza) lincolni</i> , (Aud.) Baird. Lincoln's Finch. United States from Atlantic to Pacific, and south through Mexico to Guatemala	482
369. <i>Melospiza (Melospiza) palustris</i> , (Wils.) Baird. Swamp Sparrow. Eastern United States from the Atlantic to the Missouri	483
370. <i>Peucaea aestivalis</i> , (Licht.) Cab. Bachman's Finch. Georgia	484
371. <i>Peucaea cassinii</i> , (Woodh.) Baird. San Antonio, Texas, to Los Nogales, Sonora	485
372. <i>Peucaea ruficeps</i> , (Cass.) Baird. Coast of California	486
373. <i>Embernagra rufivirgata</i> , Lawr. Valley of the Rio Grande and probably of Gila, southward. Mazatlan, Mexico	487
374. <i>Passerella iliaca</i> , (Merrem.) Sw. Fox-colored Sparrow. Eastern United States to the Mississippi.	488
375. <i>Passerella townsendii</i> , (Aud.) Nutt. Pacific coast of United States as far south as Sacramento. Fort Tejon ??	489
376. <i>Passerella schistacea</i> , Baird. Head-waters of Platte and Fort Bridger to Fort Tejon, California. Perhaps another species, <i>P. megarhynchus</i> , (App.) 490, 925	490, 925
377. <i>Calamospiza bicolor</i> , (Town.) Bon. Lark Bunting. High Central Plains to the Rocky mountains; southwesterly to valley of Mimbres and Sonora	492
378. <i>Euspiza americana</i> , (Gm.) Bon. Black-throated Bunting. United States from the Atlantic to the border of the High Central Plains	494

	Page.
379. <i>Euspiza townsendii</i> , (Aud.) Bon. Townsend's Bunting. Chester county, Pennsylvania. But one specimen known.....	495
380. <i>Guiraca (Goniaphea) ludoviciana</i> , (Linn.) Sw. Rose-breasted Grosbeak. Eastern United States to the Missouri plains; south to Guatemala.....	497
381. <i>Guiraca (Goniaphea) melanocephala</i> , Sw. Black-headed Grosbeak. High Central Plains from Yellow Stone to the Pacific; Table lands of Mexico.....	498
382. <i>Guiraca caerulea</i> , (Linn.) Sw. Blue Grosbeak. More Southern United States from Atlantic to Pacific; south to Mexico.....	499
383. <i>Cyanospiza parellina</i> , (Bon.) Baird. Northeastern Mexico to the Rio Grande.....	502
384. <i>Cyanospiza ciris</i> , (Linn.) Baird. Painted Bunting. South Atlantic and Gulf States to the Pecos river, Texas; south into Mexico.....	503
385. <i>Cyanospiza versicolor</i> , (Bon.) Baird. Northeastern Mexico, probably to the Rio Grande; Peru.....	503
386. <i>Cyanospiza amoena</i> , (Say.) Baird. Lazuli Finch. High Central Plains to the Pacific.....	504
387. <i>Cyanospiza cyanea</i> , (Linn.) Baird. Indigo Bird. Eastern United States to the Mis- souri; south to Guatemala.....	505
388. <i>Spermophila moreletii</i> , Pucheran. Rio Grande of Texas; south to Honduras.....	506
389. <i>Pyrrhuloxia sinuata</i> , Bon. Valley of the Rio Grande of Texas.....	508
390. <i>Cardinalis virginianus</i> , Bonaparte. Red Bird. More southern portions of the Unsted States to the Missouri; probably along valley of Rio Grande to Rocky mountains.....	509
391. <i>Pipilo erythrophthalmus</i> , (Linn.) Vieill. Ground Robin. Eastern United States to the Missouri river.....	512
392. <i>Pipilo oregonus</i> , Bell. Oregon Ground Robin. Coast of Oregon and Washington Territories.....	513
393. <i>Pipilo arcticus</i> , Sw. High Central Plains of Upper Missouri, Yellowstone, and Platte; Fort Bridger.....	514
394. <i>Pipilo megalonyx</i> , Baird. Southern coast of California and across through vallies of Gila and Rio Grande.....	515
395. <i>Pipilo aberti</i> , Baird. Base of Rocky mountains in New Mexico; Valley of Gila and Colorado.....	516
396. <i>Pipilo fuscus</i> , Sw. Coast region of California.....	517
397. <i>Pipilo mesoleucus</i> , Baird. Valley of Upper Rio Grande and across to the Gila river; East to Santa Caterina, New Leon.....	518
398. <i>Pipilo chlorurus</i> , (Town.) Baird. Blanding's Finch. Valley of Rio Grande and Gila. Rocky mountains north to the South Pass; south to Mexico.....	519
399. <i>Dolichonyx oryzivorus</i> , (Linn.) Sw. Boblink. Eastern United States to Fort Bridger, Utah.....	522
400. <i>Molothrus pecoris</i> , (Gm.) Sw. Cow Bird. United States from the Atlantic to Cali- fornia; Fort Bridger.....	524
401. <i>Agelaius phoeniceus</i> , (Linn.) Vieill. Swamp Blackbird. United States from Atlantic to Pacific.....	526

402. <i>Agelaius gubernator</i> , (Wagl.) Bon. Red-shouldered Blackbird. Pacific coast of United States; Colorado river	529
403. <i>Agelaius tricolor</i> , (Nutt.) Bon. Red and White-shouldered Blackbird. Coast of California; Colorado river	530
404. <i>Xanthocephalus icterocephalus</i> , (Bon.) Baird. Yellow-headed Blackbird. Western America, from Texas, Illinois, Wisconsin, and North Red river to California; south into Mexico; Greenland	531
(405.) <i>Trupialis militaris</i> , (Linn.) Bonap. Red-breasted Lark. Western coast of South America; around to Falkland Islands; perhaps in Brazil; coast of California?	533
406. <i>Sturnella magna</i> , (Linn.) Sw. Meadow Lark. Eastern United States to the High Central Plains; south to Mexico; Cuba?	535
407. <i>Sturnella neglecta</i> , Aud. Western Lark. Western America, from High Central Plains to the Pacific; east to Pembina, and perhaps to Wisconsin	537
408. <i>Icterus vulgaris</i> , (Linn.) Daudin. Troupial. Northern South America and West Indies; accidental on the southern coast of United States	542
409. <i>Icterus audubonii</i> , Giraud. Audubon's Oriole. Valley of the Lower Rio Grande of Texas, southward	542
(410.) <i>Icterus melanocephalus</i> , (Wagl.) Gray. Warm parts of Mexico	543
411. <i>Icterus parisorum</i> , Bon. Valley of the Rio Grande, south to Guatemala; in Texas, found on the Pecos	544
412. <i>Icterus wagleri</i> , Sclater. Northeastern Mexico to Rio Grande valley; south to Guatemala	545
413. <i>Icterus cucullatus</i> , Swain. Hooded Oriole. Valley of the Lower Rio Grande, southward	546
414. <i>Icterus spurius</i> , (Linn.) Bon. Orchard Oriole. United States, from the Atlantic to the High Central Plains; probably throughout Texas; south to Guatemala	547
415. <i>Icterus baltimore</i> , (Linn.) Daudin. Baltimore Oriole. From Atlantic coast to the High Central Plains, and in their borders; south to Guatemala	548
416. <i>Icterus bullockii</i> , (Sw.) Bon. Bullock's Oriole. High Central Plains to the Pacific; rare on upper Missouri; south into Mexico	549
417. <i>Scolecophagus ferrugineus</i> , (Gm.) Sw. Rusty Blackbird. From Atlantic coast to the Missouri	551
418. <i>Scolecophagus cyanocephalus</i> , (Wagl.) Cab. Brewer's Blackbird. High Central Plains to the Pacific; south to Mexico; Pembina, Minnesota	552
419. <i>Quiscalus macrourus</i> , Sw. Great-tailed Grackle. Valley of the Lower Rio Grande of Texas, southward	553
420. <i>Quiscalus major</i> , (Wils.) Vieill. Boat-tailed Grackle. Southern Atlantic and Gulf Coast; mouth of Rio Grande, Texas	555
421. <i>Quiscalus versicolor</i> , (Linn.) Vieill. Crow Blackbird. Atlantic to High Central Plains; Fort Bridger	555
422. <i>Quiscalus baritus</i> , (Linn.) Vieill. Florida coast and West Indies	556
423. <i>Corvus carnivorus</i> , Bartram. American Raven. Entire continent of North America; rare east of the Mississippi	560
424. <i>Corvus cacalotl</i> , Wagl. Colorado Raven. Colorado river of California, southward ..	563

	Page.
425. <i>Corvus cryptoleucus</i> , Couch. White-necked Crow. Valley of Rio Grande and Gila; abundant on the Llano Estacado.....	565
426. <i>Corvus americanus</i> , Aud. Common Crow. North America to the Missouri region; also on the coast of California? (Not found on the High Central Plains?).....	566
427. <i>Corvus americanus</i> , var. <i>floridanus</i> , Baird. Florida Crow. Southern peninsula of Florida.....	568
428. <i>Corvus caurinus</i> , Baird. Northwestern Fish Crow. Washington Territory and north-west coast.....	569
429. <i>Corvus ossifragus</i> , Wilson. Fish Crow. South Atlantic (and Gulf?) coast.....	571
430. <i>Picicorvus columbianus</i> , (Wils.) Bon. Clark's Crow. From Rocky mountains to Pacific; east to Fort Kearney.....	573
431. <i>Gymnokitta cyanocephala</i> , Pr. Max. Maximilian's Jay. Rocky mountains to Cascades of California and Oregon. Not on the Pacific coast.....	574
432. <i>Pica ludsonica</i> , (Sabine,) Bon. Magpie. Arctic regions of North America; the United States from the High Central Plains to the Pacific north of California.....	576
433. <i>Pica nuttalli</i> , Aud. Yellow-billed Magpie. Coast of California.....	578
434. <i>Cyanura cristata</i> , (Linn.) Sw. Blue Jay. Eastern North America; west to the Missouri.....	580
435. <i>Cyanura stelleri</i> , (Gm.) Sw. Steller's Jay. Pacific coast of North America; east to St. Mary's mission, Rocky mountains.....	581
436. <i>Cyanura macrolophus</i> , Baird. Long-crested Jay. Central line of Rocky mountains to the table lands of Mexico.....	582
437. <i>Cyanocitta californica</i> , (Vigors,) Strick. California Jay. Pacific coast from Columbia river, south; not in the interior.....	584
438. <i>Cyanocitta woodhousii</i> , Baird. Woodhouse's Jay. Central line of Rocky mountains to the table lands of Mexico.....	585
439. <i>Cyanocitta floridana</i> , (Bartram,) Bon. Florida Jay. Florida.....	586
440. <i>Cyanocitta sordida</i> , (Sw.) Baird. Mimbres region of Rocky mountains, and south to table lands of Mexico.....	587
441. <i>Cyanocitta ultramarina</i> , (Bon.) Strickl. Ultramarine Jay. South side of valley of Rio Grande, near the coast and southward.....	588
442. <i>Xanthoura luxuosa</i> , (Lesson,) Bon. Green Jay. Valley of Rio Grande of Texas, and southward.....	589
443. <i>Perisoreus canadensis</i> , (Linn.) Bon. Canada Jay. Northern America into the northern parts of United States from Atlantic to Pacific; farther south in Rocky mountains.....	590
444. <i>Psilorhynchus morio</i> , (Wagler,) Gray. Rio Grande valley of Texas, southward.....	592
445. <i>Columba (Columba) fasciata</i> , Say. Band-tailed Pigeon. Rocky mountains to Pacific coast; south to New Leon, Mexico.....	597
446. <i>Columba (Columba) flavirostris</i> , Wagl. Red-billed Dove. Lower Rio Grande.....	598
447. <i>Columba (Patagioenas) leucocephala</i> , Linn. White-headed Pigeon. Indian Key and other southern keys of Florida; not on main land; West Indies generally.....	599

448. <i>Ectopistes migratoria</i> , (Linn.) Sw. Wild Pigeon. North America to High Central Plains	600
449. <i>Zenaida amabilis</i> , Bonap. Zenaida Dove. Florida Keys; chiefly on or near Indian Key and the West Indies	602
450. <i>Melopelia leucoptera</i> , (Linn.) Bon. White-winged Dove. Valley of Rio Grande, southward; California; West Indies	603
451. <i>Zenaidura carolinensis</i> , (Linn.) Bon. Common Dove. Throughout United States, from Atlantic to Pacific; Cuba	604
452. <i>Scardafella squamosa</i> , (Temm.) Bon. Scaly Dove. South side of valley of Rio Grande, southward; perhaps <i>S. inca</i> , Bon	605
453. <i>Chamaepelia passerina</i> , (Linn.) Sw. Ground Dove. South Atlantic and Gulf coasts, Lower California; accidental near Washington, D. C.	606
454. <i>Oreopelia martinica</i> , (Gm.) Reich. Key West Pigeon. Key West, Florida, and West Indies	607
455. <i>Starnoenas cyanocephala</i> , (Linn.) Bon. Blue-headed Pigeon. West India islands; occasionally at Key West, Florida, and other southern Keys	608
456. <i>Ortalia m'c calli</i> , Baird. Chicalacca. New Mexico	611
457. <i>Meleagris gallopavo</i> , Linn. Wild Turkey. United States	615
458. <i>Meleagris mexicana</i> , Gould. Mexican Turkey. Mexico	618
459. <i>Tetrao obscurus</i> , Say. Dusky Grouse. Black Hills of Nebraska to Cascade mountains of Oregon and Washington	620
460. <i>Tetrao canadensis</i> , Linn. Spruce Partridge. Spruce forests and swamps of northern United States to the Arctic seas; west nearly to the Rocky mountains	622
461. <i>Tetrao franklinii</i> , Douglas. Franklin's Grouse. Northern Rocky mountains, and west	623
462. <i>Centrocercus urophasianus</i> , (Bon.) Sw. Cock of the Plains. Sage plains of the north-west	624
463. <i>Pediocetes phasianellus</i> , (Linn.) Baird. Sharp-tailed Grouse. Northern prairies and plains, from Wisconsin to Cascades of Oregon and Washington	626
464. <i>Cupidonia cupido</i> , (Linn.) Baird. Prairie Hen. Western prairies and plains, within the limits of the United States east of the Rocky Mountains; southeast to Calcasieu, Louisiana; east to Pocono mountains, Pennsylvania, Long Island, and eastern coast	628
465. <i>Bonasa umbellus</i> , (Linn.) Steph. Ruffed Grouse. Wooded portions of eastern United States towards the Rocky mountains	630
465.* <i>Bonasa</i> , var. <i>umbelloides</i> , (Douglas,) Baird. Gray Mountain Grouse. Northern valleys of Rocky mountains	630
466. <i>Bonasa sabinii</i> , (Douglas,) Baird. Oregon Grouse. Rocky mountains, to Pacific coast of Oregon and Washington	631
467. <i>Lagopus albus</i> , (Gm.) Aud. White Ptarmigan. Northern America; rare in the northern parts of United States	633
468. <i>Lagopus rupestris</i> , (Gm.) Leach. Ptarmigan. Arctic America	635

	Page.
469. <i>Lagopus leucurus</i> , Sw. White-tailed Ptarmigan. Northern America to the west; southward along Rocky mountains to Cochetope pass, in latitude 39°	636
470. <i>Lagopus americanus</i> , Aud. American Ptarmigan. Arctic America	637
471. <i>Ortyx virginianus</i> , (Linn.) Bon. Partridge; Quail. Eastern United States, to the High Central Plains	640
472. <i>Ortyx texanus</i> , Lawr. Southern Texas and valley of Rio Grande	641
473. <i>Oreortyx pictus</i> , (Douglas,) Baird. Plumed Partridge. Mountain ranges of California and Oregon towards the coast	642
474. <i>Lophortyx californicus</i> , (Shaw,) Bon. California Quail. Plains and lowlands of California and Oregon towards the coast; Mohave river	644
475. <i>Lophortyx gambelii</i> , Nutt. Gambel's Partridge. Upper Rio Grande and Gila, to the Colorado of California	645
476. <i>Callipepla squamata</i> , (Vigors,) Gray. Scaled or Blue Partridge. Valley of Rio Grande of Texas. Not yet detected further west. Most abundant on the high broken table lands and mezquite plains	646
477. <i>Cyrtonyx massena</i> , (Lesson,) Gould. Massena Partridge. Chiefly on the upper Rio Grande, from the high plains of the Pecos	647
478. <i>Grus americanus</i> , (Linn.) Ord. Whooping Crane. Florida and Texas; stragglers in Mississippi valley	654
479. <i>Grus canadensis</i> , (Linn.) Temm. Sand-hill Crane. Whole of western regions of United States; Florida	655
480. <i>Grus fraterculus</i> , Cassin. New Mexico	656
481. <i>Aramus giganteus</i> , (Bon.) Baird. Crying Bird. Florida and West Indies	657
482. <i>Demiegretta pealii</i> , (Bon.) Baird. Peale's Egret. Seacoast of south Florida	661
483. <i>Demiegretta rufa</i> , (Bodd.) Baird. Reddish Egret. Coast of south Florida and Gulf of Mexico to mouth of Rio Grande; Cuba	662
484. <i>Demiegretta ludoviciana</i> , (Wils.) Baird. Louisiana Heron. Coast of South Atlantic and Gulf States	663
485. <i>Garzetta candidissima</i> , (Jacquin,) Bon. Snowy Heron. Coast of middle and Gulf States, and across to California	665
486. <i>Herodias egretta</i> , (Gmel.) Gray. White Heron. Southern portions of the United States; stragglng to Massachusetts	666
486.* <i>Herodias egretta</i> , var. <i>californica</i> , Baird. Coast of southern California, and perhaps the Rio Grande of Texas	667
487. <i>Ardea herodias</i> , Linn. Great Blue Heron. Throughout the entire territory of the United States; West Indies	668
488. <i>Ardea wurdemannii</i> , Baird. Florida Heron. South Florida	669
489. <i>Audubonia occidentalis</i> , (Aud.) Bon. Great White Heron. South Florida and Cuba	670
490. <i>Florida caerulea</i> , (Linn.) Baird. Blue Heron. South Atlantic and Gulf coast to Mexico	671
491. <i>Ardetta exilis</i> , (Gmel.) Gray. Least Bittern. Throughout the United States, from Atlantic to Pacific	673
492. <i>Botaurus lentiginosus</i> , Steph. Bittern. Entire continent of North America	674

	Page.
493. <i>Butorides virescens</i> , (Linn.) Bon. Green Heron. United States generally	676
494. <i>Butorides brunnescens</i> , (Cab.) Cuba; Florida?	676
495. <i>Nyctiardea gardeni</i> , (Gmelin,) Baird. Night Heron. United States generally.	678
496. <i>Nyctherodius violaceus</i> , (Linn.) Reich. Yellow-crowned Night Heron. South Atlantic and Gulf States; South America.	679
497. <i>Tantalus loculator</i> , (Linn.) Wood Ibis. South Atlantic and Gulf States and across to the Colorado river; as far north as North Carolina and mouth of Ohio	682
498. <i>Ibis rubra</i> , (Linn.) Vieillot. Red or Scarlet Ibis. South America and West Indies. Very rare or accidental in the United State.	683
499. <i>Ibis alba</i> , (Linn.) Vieillot. White Ibis. South Atlantic and Gulf States; straggling occasionally northward	684
500. <i>Ibis (Falcinellus) ordii</i> , Bonaparte. Glossy Ibis. Found singly and at intervals over the whole United States	685
501. <i>Platalea ajaja</i> , Linn. Rosy Spoonbill. South Atlantic and Gulf States	686
502. <i>Phoenicopterus ruber</i> , Linn. Flamingo. Warm parts of America. Rare on the Florida Keys	687
503. <i>Charadrius virginicus</i> , Borek. Golden Plover. All of North America, South America, Northern Asia, Europe	690
504. <i>Aegialitis (Oxyechus) vociferus</i> , (Linn.) Cassin. Killdeer. North America to the Arctic regions; Mexico; South America.	692
505. <i>Aegialitis (Oxyechus) montanus</i> , (Townsend) Cassin. Mountain Plover. Western North America; Fort Bridger, and Fort Tejon	693
506. <i>Aegialitis (Ochthodromus) wilsonius</i> , (Ord.) Cassin. Wilson's Plover. Middle and Southern States on the Atlantic, and the Atlantic coast of South America.	693
507. <i>Aegialitis (Aegialeus) semipalmatus</i> , (Bon.) Cab. Semipalmated Plover. The whole of temperate North America. Common on the Atlantic	694
508. <i>Aegialitis (Aegialeus) melodus</i> , (Ord.) Cab. Piping Plover. Eastern coast of North America; Nebraska; Louisiana	695
509. <i>Aegialitis (Leucopoliis) nivosa</i> , Cassin. Presidio, (near San Francisco,) California	695
510. <i>Squatarola helvetica</i> , (Linn.) Cuv. Black-bellied Plover. All of North America. The seacoasts of nearly all countries of the world.	697
511. <i>Aphriza virgata</i> , (Gmelin,) Gray. Surf Bird. Pacific coast of North America? South America; Sandwich Islands	698
512. <i>Haematopus palliatus</i> , Temm. Oyster Catcher. Coast of Atlantic ocean; States on the Pacific? Florida	699
513. <i>Haematopus niger</i> , Pallas. Bachman's Oyster Catcher. Western coast of the United States; Kurile islands	700
(514)? <i>Haematopus ater</i> , Vieillot. Western coast of the United States? South America; coast of Chile	700
515. <i>Streptilas interpres</i> , (Linn.) Illig. Turnstone. Shores of the Atlantic and Pacific, throughout North America. One of the most widely diffused of birds, being found in nearly all parts of the world	701
516. <i>Streptilas melanocephalus</i> , Vigors. Black Turnstone. Western North America.	702

	Page.
517. <i>Recurvirostra americana</i> , Gm. American Avoset. All of temperate North America; Florida	703
518. <i>Himantopus nigricollis</i> , Vieillot. Black-necked Stilt. United States generally	704
519. <i>Phalaropus wilsonii</i> , Sab. Wilson's Phalarope. Entire temperate regions of north America; New Mexico	705
520. <i>Phalaropus hyperboreus</i> , (Linn.) Temm. Northern Phalarope. The whole of temperate North America; Europe; Japan; San Francisco, Cal.	706
521. <i>Phalaropus fulicarius</i> , (Linn.) Bon. Red Phalarope. Entire temperate regions of North America; Asia; Europe	707
522. <i>Philohela minor</i> , (Gm.) Gray. American Woodcock. Eastern North America	709
523. <i>Gallinago wilsonii</i> , (Temm.) Bon. English Snipe. Entire temperate regions of North America; California	710
524. <i>Macrorhamphus griseus</i> , (Gm.) Leach. Red-breasted Snipe. Entire temperate regions of North America	712
525. <i>Macrorhamphus scolopaceus</i> , (Say.) Lawrence. Entire temperate regions of North America; Washington Territory; New York	712
526. <i>Tringa (Tringa) canutus</i> , Linn. Gray Back. Eastern North America; Europe	715
527. <i>Tringa (Tringa) cooperi</i> , Baird. Long Island	716
528. <i>Tringa (Arquatella) maritima</i> , Brünnich. Purple Sandpiper. Eastern North America; Europe	717
529. <i>Tringa (Erolia) subarquata</i> , (Gould.) Temm. Curlew Sandpiper. Atlantic coast of the United States; rare; Europe; Asia; Africa	718
530. <i>Tringa (Schoeniclus) alpina</i> , var. <i>americana</i> , Cassin. Red-backed Sandpiper. Entire temperate regions of North America	719
531. <i>Tringa (Actodromas) maculata</i> , Vieill. Jack Snipe. North America generally. South America. Accidental in Europe	720
532. <i>Tringa (Actodromas) wilsonii</i> , Nuttall. Least Sandpiper. Entire temperate North America.	721
533. <i>Tringa (Actodromas) bonapartii</i> , Schlegel. North America east of the Rocky mountains	722
534. <i>Calidris arenaria</i> , (Linn.) Illiger. Sanderling. Entire temperate regions of North America; South America; Europe	723
535. <i>Ereunetes petrificatus</i> , Ill. Semipalmated Sandpiper. Entire temperate regions of North America; South America; varying much in size	724
536. <i>Micropalama himantopus</i> , (Bon.) Baird. Stilt Sandpiper. Eastern North America	726
537. <i>Symphemia semipalmata</i> , (Gm.) Hartlaub. Willet. Entire temperate regions of North America; South America	729
538. <i>Glottis floridanus</i> , Bon. Florida Greenshank. Florida	730
539. <i>Gambetta melanoleuca</i> , (Gm.) Bon. Tell-tale. Stone Snipe. Entire temperate regions of North America. Mexico	731
540. <i>Gambetta flavipes</i> , (Gm.) Bon. Yellow Legs. Eastern North America; western?	732
541. <i>Rhyacophilus solitarius</i> , (Wils.) Bon. Solitary Sandpiper. Entire temperate regions of North America. Mexico	733
542. <i>Heteroscelus brevipes</i> , (Vieill.) Baird. Wandering Tattler. Washington Territory. Islands in the Pacific. South America. Northeastern Asia. Japan	734

543. <i>Tringoides macularius</i> , (Linn.) Gray. Spotted Sandpiper. Entire temperate North America; Oregon. Accidental in Europe	735
544. <i>Philomachus pugnax</i> , (Linn.) Gray. Ruff. Northern Europe and Asia. Accidental on Long Island	737
545. <i>Actiturus bartramius</i> , (Wils.) Bon. Field Plover. Eastern North America; South America. Europe	737
546. <i>Tryngites rufescens</i> , (Vieillot,) Cab. Buff-breasted Sandpiper. All of North America; South America; Europe	739
547. <i>Limosa fedoa</i> , (Linn.) Ord. Entire temperate regions of North America. South America	740
548. <i>Limosa hudsonica</i> , (Lath.) Sw. Northern and eastern North America; New Jersey .	741
549. <i>Numenius (Numenius) longirostris</i> , Wilson. Long-billed Curlew. The entire temperate regions of North America. Perhaps two or more species included	743
550. <i>Numenius (Phaeopus) hudsonicus</i> , Latham. Hudsonian Curlew. Atlantic and Pacific coasts of North America. California	744
551. <i>Numenius (Phaeopus) borealis</i> , (Forst.) Latham. Esquimaux Curlew. Eastern and northern North America	744
552. <i>Rallus elegans</i> , Aud. Marsh Hen. Middle and Southern States on the Atlantic Ocean; California	746
553. <i>Rallus crepitans</i> , Gm. Clapper Rail. Middle and southern coast of the States on the Atlantic Ocean; South America	747
554. <i>Rallus virginianus</i> , Linn. Virginia Rail. The entire temperate regions of North America; New Mexico, California, Oregon	748
555. <i>Porzana (Porzana) carolina</i> , Vieill. Common Rail. Entire temperate regions of North America	749
556. <i>Porzana (Creciscus) jamaicensis</i> , (Gm.) Little Black Rail. Middle and Southern States on the Atlantic Ocean	749
557. <i>Porzana (Coturnicops) noveboracensis</i> , (Gm.) Yellow Rail. Eastern North America .	750
558. <i>Crex pratensis</i> , Bechst. Corn-crake. Europe; Greenland. Accidental on the Atlantic coast of the United States	751
559. <i>Fulica americana</i> , Gmelin. Coot. Entire temperate regions of North America .	751
560. <i>Gallinula (Gallinula) galeata</i> , (Licht.) Bon. Florida Gallinule. Southern countries of North America; accidental in Middle and Northern States	752
561. <i>Gallinula (Porphyrula) martinica</i> , (Linn.) Lath. Purple Gallinule. Southern States of North America, Louisiana, Florida. Accidental in the middle and northern United States	753
561. <i>Cygnus americanus</i> , Sharpless. American Swan. Continent of North America . .	758
562. <i>Cygnus buccinator</i> , Rich. Trumpeter Swan. Western America, from the Mississippi valley to the Pacific	758
563. <i>Anser (Chen) hyperboreus</i> , Pallas. Snow Goose. Whole of North America. Perhaps a second species, (<i>A. albatus</i>)	760

	Page.
564. <i>Anser (Chen) caerulescens</i> , Linn. White-headed Goose. North America.....	761
565. <i>Anser (Anser) gambelii</i> , Hartlaub. White Fronted Goose. Whole of North America	761
566. <i>Anser frontalis</i> , Baird. Interior of North America.....	762
567. <i>Bernicla (Leucoblepharon) canadensis</i> , (Linn.) Boie. Canada Goose. Whole of North America. Accidental in Europe	764
568. <i>Bernicla (Leucoblepharon) leucopareia</i> , (Brandt,) Cassin. White-cheeked Goose. West coast of America. Perhaps mixed with <i>B. occidentalis</i>	765
569. <i>Bernicla (Leucoblepharon) hutchinsii</i> , (Rich.) Bonap. Huetchin's Goose. Northern and western regions of North America	765
570. <i>Bernicla (Bernicla) brenta</i> , Steph. Brant. Eastern or Atlantic coast of North America; Europe. Not yet observed on the Pacific side of the continent	767
571. <i>Bernicla (Bernicla) nigricans</i> , (Lawr.) Cassin. Black Brant. Pacific coast of North America. Very rare on the Atlantic coast.....	767
572. <i>Bernicla (Leucopareia) leucopsis</i> , (Linn.) Barnacle Goose. Very doubtful as an inhabitant of North America	768
573. <i>Chloephaya canagica</i> , (Sewast.) Bon. Painted Goose. Aleutian Islands.....	768
574. <i>Dendrocygna autumnalis</i> , (Linn.) Eyton. Long-legged Duck. Valley of Rio Grande, Texas; also in South America and West Indies.....	770
575. <i>Dendrocygna fulva</i> , (Gmelin,) Burm. Fort Tejon, California, and south into Brazil.	770
576. <i>Anas boschas</i> , Linn. Mallard. Entire continent of North America and greater part of Old World.....	774
577. <i>Anas obscura</i> , Gm. Black Duck. Atlantic region of North America. Not yet detected on the Pacific, nor in Europe	775
578. <i>Dafila acuta</i> , (Linn.) Jenyns. Sprig-tail; Pin-tail. Whole of North America and Europe	776
579. <i>Nettion carolinensis</i> , (Gm.) Baird. Green-winged Teal. Whole of North America; accidental in Europe.....	777
580. <i>Nettion crecca</i> , (Linn.) Kaup. English Teal. Europe; accidental on the eastern coast of the United States.....	778
581. <i>Querquedula discors</i> , (Linn.) Steph. Blue-winged Teal. Eastern North America to Rocky mountains. Not yet found on the Pacific coast, nor in Europe	779
582. <i>Querquedula cyanopectera</i> , (Vieill.) Baird. Red-breasted Teal. Rocky mountains to Pacific; accidental in Louisiana; spread over most of western South America.....	780
583. <i>Spatula clypeata</i> , (Linn.) Boie. Shoveller. Continent of North America; abundant in Europe.....	781
584. <i>Chaulelasmus streperus</i> , (Linn.) Gray. Gadwall. North America generally, and Europe	882
585. <i>Mareca americana</i> , (Gm.) Stephens. Baldpate; American Widgeon. Continent of North America; accidental in Europe.....	783

	Page.
586. <i>Mareca penelope</i> , (Linn.) Bon. English Widgeon. Old World; accidental on the Atlantic coast of United States; Greenland	784
587. <i>Aix sponsa</i> , (Linn.) Boie. Summer Duck. Continent of North America	785
588. <i>Fulix marila</i> , (Linn.) Baird. Big Black-head. Whole of North America and Europe	791
589. <i>Fulix affinis</i> , (Forster,) Baird. Little Black-head. Whole of North America; accidental in Europe	791
590. <i>Fulix collaris</i> , (Donovan,) Baird. Ring-necked Duck. Whole of North America; accidental in Europe	792
591. <i>Aythya americana</i> , (Eyton,) Bon. Red-head. Whole of North America	793
592. <i>Aythya vallisneria</i> , (Wils.) Bon. Canvas-back. Whole of North America	794
593. <i>Bucephala americana</i> , (Bon.) Baird. Golden Eye. Whole of North America	796
594. <i>Bucephala islandica</i> , (Gm.) Baird. Barrow's Golden Eye. Iceland and northern parts of America; in winter, not rare on the St. Lawrence	796
595. <i>Bucephala albeola</i> , (Linn.) Baird. Butter Ball. Whole of North America	797
596. <i>Histrionicus torquatus</i> , (Linn.) Bon. Harlequin Duck. Northern seacoast of northern hemisphere	799
597. <i>Harelda glacialis</i> , (Linn.) Leach. South Southerly. Near both coasts of North America; Europe	800
598. <i>Polysticta stelleri</i> , (Pallas,) Eyton. Steller's Duck. Northeastern Asia; accidental (?) on northwest coast of America, and in Europe	801
599. <i>Lampronetta fischeri</i> , Brandt. Spectacled Eider. Norton sound, Russian America, 63½ N. L	803
600. <i>Camptolaemus labradorius</i> , (Gm.) Gray. Labrador Duck. Northeastern coast of North America	803
601. <i>Melanetta velvetina</i> , (Cassin,) Baird. Velvet Duck. Near both coasts of North America, to the north; perhaps <i>M. carbo</i> , of Pallas	805
602. <i>Pelionetta perspicillata</i> , (Linn.) Kaup. Surf Duck. On and near seacoast of North America; quite far south in winter; accidental in Europe	806
603. <i>Pelionetta trowbridgii</i> , Baird. Long-billed Scoter. Coast of southern California in winter	806
604. <i>Oidemia americana</i> , Swains. Scoter. Seacoast of North America	807
605. <i>Oidemia (Pelionetta) bimaculata</i> , Baird. Huron Scoter. Lake Huron and adjacent waters in fall and winter	808
606. <i>Somateria mollissima</i> , (Linn.) Leach. Eider Duck. Atlantic and Arctic coasts of northern hemisphere; Pacific coast, N. A	809
607. <i>Somateria V. nigra</i> , Gray. Pacific Eider. Kotzbue sound, N. W. coast of America	810
608. <i>Somateria spectabilis</i> , (Linn.) Leach. King Eider. Arctic regions of northern hemisphere; Pacific coast	810
609. <i>Erismatura rubida</i> , (Wils.) Bon. Ruddy Duck. Whole of North America; abundant throughout the interior	811
610. <i>Erismatura dominica</i> , (Linn.) Eyton. Black Masked Duck. West Indies; accidental on Lake Champlain	811

	Page.
611. <i>Mergus americanus</i> , Cass. Sheldrake. Whole of North America	813
612. <i>Mergus serrator</i> , Linn. Red-breasted Merganser. Whole of North America and Europe.	814
613. <i>Lophodytes cucullatus</i> , (Linn.) Reich. Hooded Merganser. Whole of North America	816
614. <i>Mergellus albellus</i> , (Linn.) Selby. Smew. Northern parts of Old World; very accidental in America	817
615. <i>Pelecanus (Cyrtopelecanus) erythrorhynchus</i> , Gmelin. American Pelican. North America generally, on both shores and in the interior	868
616. <i>Pelecanus (Onocrotalus) fuscus</i> , Linn. Brown Pelican. South Atlantic and Gulf coasts of the United States; coast of California	870
617. <i>Sula (Sula) bassana</i> , Briss. Common Gannet: Solan Goose. Atlantic coast of North America and Europe	871
618. <i>Sula (Dysporus) fiber</i> , (Linn.) Booby Gannet. South Atlantic and Gulf coast of United States	872
619. <i>Tachypetes aquilus</i> , (Linn.) Vieillot. Frigate Pelican: Man-of-war Bird. South Atlantic and Gulf coasts of United States and southern coast of California. Generally distributed in tropical regions	873
620. <i>Graculus (Phalacrocorax) carbo</i> , (Linn.) Gray. Common Cormorant. Atlantic coast of North America to New Jersey, (in winter) and Europe	876
621. <i>Graculus (Phalacrocorax) perspicillatus</i> , (Pallas,) Lawrence. Pallas's Cormorant. Russian America	877
622. <i>Graculus (Phalacrocorax) cincinnatus</i> , (Brandt,) Gray. Tufted Cormorant. Sitka, Russian America	877
623. <i>Graculus dilophus</i> , (Sw.) Gray. Double Crested Cormorant. Arctic North America; south to Carolina and California in winter.	877
624. <i>Graculus floridanus</i> , (Aud.) Bon. Florida Cormorant. South Atlantic and Gulf coasts of United States and some distance inland	879
625. <i>Graculus mexicanus</i> , (Brandt,) Bon. Mexican Cormorant. Coast of Texas and Rio Grande; Cuba	879
626. <i>Graculus (Urile) penicillatus</i> , (Brandt,) Bonap. Brandt's Cormorant. West coast of North America	880
627. <i>Graculus (Urile) violaceus</i> , (Gmelin,) Gray. Violet Green Cormorant. West coast of North America	881
628. <i>Plotus ankinga</i> , Linn. Snake Bird: Water Turkey. Fresh waters of South Atlantic and Gulf States	883
629. <i>Phaeton flavirostris</i> , Brandt. Yellow-billed Tropic Bird. Tropical regions of the Atlantic seas	855
630. <i>Diomedea (Diomedea) exulans</i> , Linn. The Wandering Albatross. Pacific ocean	821
631. <i>Diomedea (Phoebastria) brachyura</i> , Temm. The Short-tailed Albatross. North Pacific; coasts of California and Oregon	822
632. <i>Diomedea (Thalassarche) chlororhyncha</i> , Gmel. The Yellow-nosed Albatross. Pacific ocean; coast of Oregon	822
633. <i>Diomedea (Phoebetria) fuliginosa</i> , Gmel. The Sooty Albatross. Pacific coasts of California and Oregon.	823

	Page.
634. <i>Procellaria (Ossifragus) gigantea</i> , Gmel. The Gigantic Fulmar. Pacific ocean, off Columbia river	825
635. <i>Procellaria (Fulmarus) glacialis</i> , Linn. The Fulmar Petrel. Northern Atlantic . .	825
636. <i>Procellaria pacifica</i> , Aud. The Pacific Fulmar. Pacific coasts of North America .	826
637. <i>Procellaria (Thalassoica) tenuirostris</i> , Aud. The Slender-billed Fulmar. Pacific coast; Columbia river	826
638. <i>Procellaria (Aestrelata) meridionalis</i> , Lawrence. The Tropical Fulmar. Atlantic coast, from Florida to New York	827
639. <i>Daption capensis</i> , (Linn.) Steph. The Pintado Petrel. Off the coast of California .	828
640. <i>Thalassidroma (Oceanodroma) furcata</i> , (Gmel.) Gould. Fork-tailed Petrel. Coasts of Oregon and Russian America	829
641. <i>Thalassidroma (Oceanodroma) hornbyi</i> , Gray. Hornby's Petrel. Northwest coast of America	829
642. <i>Thalassidroma leachii</i> , Temm. Leach's Petrel. Atlantic coast, from Massachusetts to Baffin's bay	830
643. <i>Thalassidroma melanota</i> , Bon. The Black Stormy Petrel. Coast of California . . .	830
644. <i>Thalassidroma (Oceanites) wilsoni</i> , Bon. Wilson's Stormy Petrel. Off the Atlantic coast, from the Gulf of Mexico to Baffin's bay	831
645. <i>Thalassidroma (Procellaria) pelagica</i> , (Linn.) Bon. Mother Cary's Chicken. Atlantic ocean, banks of Newfoundland	831
646. <i>Fregatta lawrencii</i> , Bonap. The Black and White Stormy Petrel. Florida coast . .	832
647. <i>Puffinus (Ardenna) major</i> , Faber. The Greater Shearwater. Atlantic ocean, Florida coast to the Gulf of St. Lawrence	833
648. <i>Puffinus (Nectris) fuliginosus</i> , Strick. The Sooty Shearwater. Atlantic coast of the Northern States. Banks of Newfoundland	834
649. <i>Puffinus anglorum</i> , Temm. The Mank's Shearwater. Coast of New Jersey to Labrador	834
650. <i>Puffinus obscurus</i> , (Gmel.) Lath. The Ducky Shearwater. Southern coast of the United States; Gulf of Mexico	835
651. <i>Puffinus (Adamastor) cinereus</i> , Gmel. The Cinereous Petrel. Pacific ocean, off the California coast	835
652. <i>Stercorarius catarractes</i> , (Linn.) Temm. The Common Skua. Coast of California .	838
653. <i>Stercorarius pomarinus</i> , Temm. The Pomarine Skua. Labrador; as far south as New York in winter	838
654. <i>Stercorarius parasiticus</i> , (Linn.) Temm. The Arctic Skua. Arctic America; coast of United States from New York, northwest	839
655. <i>Stercorarius cephus</i> , (Brünn.) Buffon's Skua. Arctic seacoasts of America; Baffin's bay	840
656. <i>Larus glaucus</i> , Brünn. The Glaucous Gull. Arctic seas, Labrador, New York in winter, rarely	842
657. <i>Larus glaucescens</i> , Licht. The Glaucous-winged Gull. Northwest coast of North America	842
658. <i>Larus leucopterus</i> , Faber. The White-winged Gull. Arctic seas; Baffin's Bay; Labrador	843
659. <i>Larus chalcopterus</i> , (Bruch.) Lawr. The Gray-winged Gull. American coast of Behring's Straits, and Greenland	843

	Page.
660. <i>Larus marinus</i> , Linn. The Great Black-backed Gull. North Atlantic, Labrador; as far south as Florida in winter	844
661. <i>Larus argentatus</i> , Brünn. The Herring Gull. Atlantic coast, from Texas to Newfoundland; western States, Ohio and Mississippi rivers	844
662. <i>Larus occidentalis</i> , Aud. The Western Gull. Northwest coast of America	845
663. <i>Larus californicus</i> , Lawr. The California Gull. West coast of North America	846
664. <i>Larus delawarensis</i> , Ord. The Ring-billed Gull. Arctic America; Texas to Labrador; western rivers; northwest coast	846
665. <i>Larus suckleyi</i> , Lawr. Suckley's Gull. Pacific coast, Puget's Sound	848
666. <i>Blasippus heermanni</i> , (Cass.) Bon. The White-headed Gull. Coast of California	848
667. <i>Chroicocephalus atricilla</i> , Linn. The Laughing Gull. Texas to Massachusetts	850
668. <i>Chroicocephalus franklinii</i> , (Rich.) Bruch. Franklin's Rosy Gull. Missouri river; interior of fur countries	951
669. <i>Chroicocephalus cucullatus</i> , (Licht.) Bruch. The Hooded Gull. Panama, Louisiana	851
670. <i>Chroicocephalus philadelphia</i> , (Ord.) Lawrence. Bonaparte's Gull. Texas to Nova Scotia; Mississippi river; fur countries, Pacific coast of North America	852
671. <i>Chroicocephalus minutus</i> , (Pallas,) Bruch. The Little Gull. Arctic America,? Europe	853
672. <i>Rissa tridactyla</i> , (Linn.) Bonap. The Kittiwake Gull. Fur countries, Labrador; southern coast in winter	854
673. <i>Rissa septentrionalis</i> , Lawr. The North Pacific Kittiwake. Pacific coast of North America, Puget's Sound	854
674. <i>Rissa brevirostris</i> , Brandt. The Short-billed Kittiwake. Northwest coast of North America	855
675. <i>Rissa nivea</i> , (Pallas,) Bruch. The Yellow-billed Gull. Russian America	855
676. <i>Pagophila eburnea</i> , (Gm.) Kaup. The Ivory Gull. Coasts of arctic America, Labrador, Newfoundland	856
677. <i>Pagophila brachytarsi</i> , Hollb. The Short-legged Gull. Greenland	856
678. <i>Rhodostethia rosea</i> , (Jard.) The Wedge-tailed Gull. Arctic seas	857
679. <i>Creagrus furcatus</i> , (Neboux,) Bon. The swallow-tailed Gull. California	857
680. <i>Xema sabinii</i> , (Sabine,) Bon. The Fork-tailed Gull. Nova Scotia, northward; arctic seas	857
681. <i>Sterna aranea</i> , Wils. The Marsh Tern. Coast of the United States, as far north as Connecticut	859
682. <i>Sterna caspia</i> , Pallas. The Caspian Tern. Coast of New Jersey, northward	859
683. <i>Sterna regia</i> , Gambel. The Royal Tern. Atlantic coast of the southern and middle States, and California	859
684. <i>Sterna elegans</i> , Gambel. The Elegant Tern. Coast of South California	860
685. <i>Sterna acyflavida</i> , Cabot. Cabot's Tern. Texas to Florida	860
686. <i>Sterna havelli</i> , Aud. Havell's Tern. Texas to South Carolina	861
687. <i>Sterna trudeaui</i> , Aud. Trudeau's Tern. Coasts of New Jersey and Long Island	861
688. <i>Sterna fuliginosa</i> , Gmelin. The Sooty Tern. Texas to Florida	861
689. <i>Sterna wilsoni</i> , Bon. Wilson's Tern. Texas to Labrador	861

	Page.
690. <i>Sterna macroura</i> , Naum. The Arctic Tern. Coast of the New England States to the Arctic seas; fur countries	862
691. <i>Sterna forsteri</i> , Nutt. Forster's Tern. Louisiana to Florida; New York; fur countries, and California	862
692. <i>Sterna paradisæa</i> , Brünn. The Roseate Tern. Florida to New York	863
693. <i>Sterna pikei</i> , Lawr. The Slender-billed Tern. Coast of California	863
694. <i>Sterna frenata</i> , Gambel. The Least Tern. Texas to Labrador; western rivers ..	864
695. <i>Hydrochelidon plumbea</i> , (Wils.) The Short-tailed Tern. Texas to the New England States; Mississippi rivers and tributaries; fur countries	864
696. <i>Anous stolidus</i> , (Linn.) Leach. Noddy Tern. Texas to Florida	865
697. <i>Rhynchops nigra</i> , Linn. Black Skimmer. From Texas to New Jersey	866
698. <i>Colymbus torquatus</i> , Brünnich. Loon; Northern Diver. Northern regions of northern hemisphere	888
699. <i>Colymbus arcticus</i> , Linn. Black-throated Diver. Arctic regions of northern hemisphere	888
700. <i>Colymbus pacificus</i> , Lawrence. Pacific Diver. Pacific coast of United States ...	889
701. <i>Colymbus septentrionalis</i> , Linn. Red-throated Diver. Northern hemisphere ...	890
702. <i>Podiceps griseigena</i> , (Bodd.) Gray. Red-necked Grebe. Northern hemisphere. American bird perhaps distinct	892
703. <i>Podiceps cristatus</i> , (Linn.) Lath. Crested Grebe. Northern hemisphere. Perhaps a second species on Pacific coast, (<i>P. cooperi</i> .)	893
704. <i>Podiceps occidentalis</i> , Lawr. Western Grebe. Pacific coast of United States ...	894
705. <i>Podiceps darkii</i> , Lawr. Clark's Grebe. New Mexico and California	895
706. <i>Podiceps cornutus</i> , (Gm.) Latham. Horned Grebe. North America	895
707. <i>Podiceps californicus</i> , Heermann. California Grebe. Western North America ..	896
708. <i>Podiceps auritus</i> , (Linn.) Lath. Eared Grebe. Northern Europe; accidental (?) in North America	897
709. <i>Podylimbus podiceps</i> , (Linn.) Lawr. North America	898
710. <i>Alca (Chenalopez) impennis</i> , Linn. Great Auk. Arctic Seas; south to Newfoundland.	900
711. <i>Alca (Utamania) torda</i> , Linn. Razor-billed Auk. Arctic Seas; south to New Jersey.	901
713. <i>Mormon (Lunda) cirrhata</i> , (Pallas,) Bon. Tufted Puffin. North Pacific Ocean; Arctic Seas, south to California	902
713. <i>Mormon (Fratercula) corniculata</i> , Naumann. The Horned Puffin. Northern Seas.	902
814. <i>Mormon (Fratercula) glacialis</i> , Leach. Sea Parrot; Puffin. Northern Seas	903
715. <i>Mormon (Fratercula) arctica</i> , (Linn,) Illiger. Puffin. North Atlantic	903
716. <i>Sagmatorrhina labradoria</i> , (Gmelin,) Cassin. North Pacific	904
717. <i>Cerorhina monocerata</i> , (Pallas,) Cassin. North Pacific Coasts; south to California.	905
718. <i>Cerorhina suckleyi</i> , Cassin. Puget Sound	906
719. <i>Phaleris (Simorhynchus) cristatellus</i> , (Pallas,) Bon. Coasts of North Pacific	906
720. <i>Phaleris (Tylorhampus) tetracula</i> , (Pallas,) Stephens. Coasts of North Pacific ...	907
721. <i>Phaleris (Tylorhampus) camtschatica</i> , (Lepechin,) Cassin. Coasts of North America	908
722. <i>Phaleris (Ciceronia) microceros</i> , Brandt. Coasts of North Pacific	908
723. <i>Phaleris (Ciceronia) pusillus</i> , Pallas. Asiatic coast of North Pacific; American do.	909
724. <i>Ptychorhampus aleuticus</i> , (Pallas,) Brandt. American Coast of North Pacific, south to California	910

LIST OF SPECIES.

LV

	Page.
725. <i>Ombria psittacula</i> , (Pallas,) Eschsch. Coast of North Pacific	910
726. <i>Uria (Uria) grylle</i> , (Linnaeus,) Latham. Guillemot. Arctic and Northern Seas. South to New Jersey	911
727. <i>Uria (Uria) columba</i> , (Pallas,) Cassin. Coasts of North Pacific	912
728. <i>Uria (Uria) carbo</i> , Pallas. Northern coasts of North Pacific	913
729. <i>Uria (Cataractes) lomvia</i> , Brännich. Foolish Guillemot. Arctic Seas and North Atlantic	913
730. <i>Uria (Cataractes) ringvia</i> , Brännich. Murre. Arctic and Northern Seas; south to California	914
731. <i>Uria (Cataractes) arra</i> , Pallas. Thick-billed Guillemot. Arctic Seas and North Atlantic; south to New Jersey	915
732. <i>Brachyrhamphus (Apobapton) marmoratus</i> , (Gmelin,) Brandt. Marbled Guillemot. American coasts of North Pacific. California	915
733. <i>Brachyrhamphus (Apobapton) wrangelii</i> , Brandt. Aleutian Islands	917
734. <i>Brachyrhamphus (Apobapton) brachypterus</i> , Brandt. Unalaska	917
735. <i>Brachyrhamphus (Apobapton) kittlitaii</i> , Brandt. Kamtschatka; North Pacific	917
736. <i>Brachyrhamphus (Synthliboramphus) antiquus</i> , (Gmelin,) Brandt. Gray-headed Auk. Northwestern coast, America	916
737. <i>Brachyrhamphus (Apobapton) temminckii</i> , Brandt. Coast of North Pacific; Puget's sound	916
738. <i>Mergulus alle</i> , (Linnaeus,) Vieillot. Sea Dove. Dove-kie. Arctic Seas and North Atlantic; south to New Jersey	918

The following birds are enumerated in the preceding list which are not legitimately entitled to a place in the fauna of North America, (exclusive of Mexico.) Some of them have been described in the report for the purpose of comparison with closely allied species of the United States; others are mentioned because introduced by previous writers, though probably on erroneous data. Future investigations will doubtless result in the removal of others from the list now retained there:

- No. 4. *Cathartes burrovianus*, Cassin. Mexico.
 63. *Rhynchopsitta pachyrhyncha*, Bon. Mexico.
 71. **Campephilus imperialis*, Gray. Mexico.
Dryotomus lineatus.¹ Mexico; South America.
 129. **Tyrannus melancholicus*, Vieill. South Mexico.
 132. *Myiarchus cooperi*, Baird. Mexico.
 171. **Geothlypis velatus*, Cab. Southern West Indies and South America.
 216. *Cardellina rubra*, Bonap. Mexico.
 219. *Setophaga miniata*, Sw. Mexico.
 224. *Euphonia elegantissima*, Gray. Mexico.
 244. **Vireo virescens*, Vieill.¹ South America.(?)
 292. *Parus meridionalis*, Scater. South Mexico.
 297. *Psaltriparus melanotus*, Bon. Guatemala.
 298. *Carpodacus hæmorrhous*, Scater. Mexico.
 311. **Chrysomitris stanleyi*, Bonap. South America.

¹ No North American specimens seen.

- 312. **Chrysomitris yarrelli*, *Bonap.* South America.
- 350. *Junco cinereus*, *Scater.* Mexico.
- 405. *Trupialis militaris*, *Bon.* South America.
- 408. **Icterus vulgaris*, *Daud.*(¹) South America.
- 410. *Icterus melanocephalus*, *Gray.* Mexico.
- 494. *Butorides brunnescens*, *Baird.* Cuba.
- 498. **Ibis rubra*, *L.*(¹) West Indies, Cuba, and South America.
- 514. **Haematopus ater*, *Vicill.* South America.

Total of species, 23 ; of which one is not mentioned in the list, leaving 22. Of the 23 species, nine marked with an asterisk are given by Mr. Audubon.

The following species, claiming to be actually inhabitants of North America, have not been described from the specimens, none having been procurable for the purpose. Of several of them no specimens are known in any collection :

<i>Haliaeetus washingtonii.</i>	<i>Thalassidroma hornbyi.</i>
<i>Regulus cuvierii.</i>	<i>melania.</i>
<i>Dendroica montana.</i>	<i>Larus chalcopterus.</i>
<i>carbonata.</i>	<i>Rissa brevirostris.</i>
<i>Myiodioides minutus.</i>	<i>nivea.</i>
<i>bonapartii.</i>	<i>Pagophila brachytarsi.</i>
<i>Ægiothus canescens.</i>	<i>Rhodostethia rosea.</i>
<i>Leucosticte griseinucha.</i>	<i>Creagrus furcatus.</i>
<i>arctous.</i>	<i>Xema sabinni.</i>
<i>Lagopus americanus.</i>	<i>Chroicocephalus minutus.</i> (¹)
<i>Chloephaga canagica.</i>	<i>Podiceps auritus.</i> (¹)
<i>Polysticta stelleri.</i> (¹)	<i>Sagmatorhina labradoria.</i>
<i>Oidemia bimaculata.</i>	<i>Brachyrhamphus kittlitzii.</i>
<i>Somateria v-nigra.</i>	<i>wrangellii.</i>
<i>Graculus perspicillatus.</i>	<i>brachypterus.</i>
<i>cinnatus.</i>	Total—31 species.

The following species are probably accidental visitors only, and are not yet entitled to a permanent place in our fauna :

<i>Milvulus tyrannus.</i>	<i>Heliornis surinamensis.</i> (²)
<i>Saxicola œnanthe.</i>	<i>Mareca penelope.</i>
<i>Chrysomitris magellanicus.</i>	<i>Nettion crecca.</i>
<i>Philomachus pugnax.</i>	<i>Erismatura dominica.</i>
<i>Crex pratensis.</i>	<i>Mergellus albellus.</i> —Total, 10 species.

SUMMARY.

Species enumerated in the list	738
Of these, extralimital	22
 Total of North American species(²)	716
North American birds given by Wilson in 1814	283
North American birds given by Bonaparte, 1838	471
North American birds given by Audubon in 1844	506

¹ No North American specimen seen.

² Not enumerated in the list.

³ Of these no specimens at all, of 28 species, were to be procured in this country for examination, and extralimital ones only of 8 others. Many supposed species are referred to in different parts of the report; some of which may prove genuine.

